

From DEPARTMENT OF LABORATORY MEDICINE

Karolinska Institutet, Stockholm, Sweden

**IMPROVING TREATMENT OUTCOMES
FOR PATIENTS WITH PULMONARY
TUBERCULOSIS IN TANZANIA:
Host and Pathogen factors**

Tumaini Joseph Nagu



**Karolinska
Institutet**

Stockholm 2017

All previously published papers were reproduced with permission from the publisher.

Published by Karolinska Institutet

Printed by Eprint AB 2017

© Tumaini Joseph Nagu, 2017

ISBN 978-91-7676-863-1

Improving treatment outcomes for patients with pulmonary tuberculosis in Tanzania: Host and Pathogen factors

THESIS FOR DOCTORAL DEGREE (Ph.D.)

Wangari hall, Tomtebodavägen 18 A, Karolinska Institutet - Solna, Stockholm

Friday the 1st of December 2017 at 09.00 a.m.

By

Tumaini Joseph Nagu

Principal Supervisor:

Professor Markus Maeurer
Karolinska Institutet
Department of Laboratory Medicine
Division of Therapeutic Medicine

Co-supervisor(s):

Professor Ferdinand Mugusi
Muhimbili University of Health and Allied
Sciences
Department of Internal Medicine

Professor Mecky I Matee
Muhimbili University of Health and Allied
Sciences
Department of Microbiology and Immunology

Doctor Rebecca Axelsson-Robertson
Karolinska Institutet
Department of Laboratory Medicine
Division of Therapeutic Immunology

Opponent:

Associate Professor Maria Lerm
Linköping University
Department of Clinical and Experimental
Medicine
Division of Microbiology and Molecular Medicine

Examination Board:

Associate Professor Carl-Johan Treutiger
Karolinska Institutet
Department of Medicine, Huddinge
Centre for infectious Medicine

Associate Professor Annelie Tjernlund
Karolinska Institutet
Department of Medicine, Solna

Professor Rune Andersson
Gothenburg University
Department of Infectious Diseases
Institute of Biomedicine, Sahlgrenska Academy

“Life is like riding a bicycle. To keep your balance, you must keep moving.”

Albert Einstein

To the loving memory of my father

ABSTRACT

Tuberculosis (TB) causes more than 1.2 million deaths each year globally. Yet, survivors of tuberculosis are left with long term sequelae of chronic inflammatory responses with consequent reduced quality of life. Biomarkers of monitoring tuberculosis disease activity, clinical response and mortality might assist in reducing mortality and ultimately in improving quality of life post tuberculosis disease.

The general aim of the thesis was to describe mortality among patients treated for a first episode of tuberculosis in Tanzania in relation to selected known pathogen related predictors of poor tuberculosis outcomes, namely HIV and anti-tuberculosis drug resistance. We also explored the role of cytokines as part of the host immune response, in relation to mortality and in modulating lung damage.

In Paper I: We show an overall mortality of 3.4 – 9.3% among patients treated for tuberculosis in Tanzania, and that TB/HIV patients were at a higher risk of death compared to TB mono-infected patients. The best benefits of antiretroviral therapy (ART) in reducing mortality for TB/HIV co-infected patients occurred when ART was initiated after 14 days of anti-tuberculosis therapy (ATT). We also observed (**Paper II**) that among 861 patients with tuberculosis, those with isoniazid resistance without concomitant rifampicin resistance had an increased risk of unsuccessful treatment outcome (death or treatment failure or loss from follow up - combined). In **Paper III**, we observed that patients who exhibited, in cell-based assays, higher interferon gamma (IFN- γ) responses against cytomegalovirus (CMV), Epstein Barr virus (EBV) or *Mycobacterium tuberculosis* ESAT-6 antigens at the time of TB diagnosis had a survival benefit following treatment for TB. However, IFN- γ responses to some viral antigens (H5N1 and HSV-1) as well as other mycobacterial antigens (Ag85A, Rv2958c, Rv0447c) were not significantly different between patients who survived and those who died. In **Paper IV**, we observed very high levels of interleukin 6 (IL-6) in serum from 234 patients with pulmonary tuberculosis, as compared to levels of IL-6 in serum from seven healthy controls. Other cytokines (IFN- γ , TNF- α , IL-2, IL-10, IL-17A and IL-21) were also analyzed at the time of TB diagnosis. Unlike with the other cytokines which returned to pretreatment levels or below following ATT, IL-6 levels at end of ATT were significantly higher than their corresponding pre-treatment levels. In addition, we also found that higher IL-6 levels at TB diagnosis correlated with survival of patients with pulmonary tuberculosis, as well as with severe lung injury, defined by chest x-ray score more than 80 at diagnosis.

In conclusion, between 3 and 9 out of 100 Tanzanian patients treated for first time tuberculosis will die during treatment. Well planned ART, appropriate clinical monitoring and timely addressing of background isoniazid resistance are essential to improve treatment outcomes. While anti CMV and EBV immune responses may serve to stratify mortality risk, adjunct therapy with IL-6 may serve as a possible target in reducing lung damage and consequently aid to improve quality of life following *Mycobacterium tuberculosis* disease in the future.

LIST OF SCIENTIFIC PAPERS

- I. **Nagu TJ**, Aboud S, Mwiru R, Matee MI, Rao M, Fawzi WW, Zumla A, Maeurer MJ, Mugusi F. Tuberculosis associated mortality in a prospective cohort in Sub Saharan Africa: Association with HIV and antiretroviral therapy. **Int J Infect Dis.** 2017; 56:39-44.
- II. **Nagu TJ**, Aboud S, Matee MI, Maeurer MJ, Fawzi WW, Mugusi F. Effects of isoniazid resistance on TB treatment outcomes under programmatic conditions in a high-TB and -HIV setting: a prospective multicentre study. **J Antimicrob Chemother.** 2017;72(3):876-881.
- III. **Nagu T**, Aboud S, Rao M, Matee M, Axelsson R, Valentini D, Mugusi F, Zumla A, Maeurer M. Strong anti-Epstein Barr virus (EBV) or cytomegalovirus (CMV) cellular immune responses predict survival and a favourable response to anti-tuberculosis therapy. **Int J Infect Dis.** 2017;56:136-139
- IV. **Nagu TJ**, Rao M, Axelsson-Robertson R, Aboud S, Matee M, Fundikira LS, Nkumbih ZF, Poiret T, Valentini D, Mugusi F, Zumla A, Maeurer M. Cytokine Biomarkers in Tanzanian patients with Pulmonary TB - a prospective longitudinal cohort study. **(Manuscript)**

CONTENTS

1	INTRODUCTION.....	7
1.1	Tuberculosis.....	7
1.1.1	Etiology and pathogenesis	7
1.1.2	Global epidemiology of tuberculosis.....	9
1.1.3	Clinical manifestation of tuberculosis	11
1.1.4	Diagnosis of tuberculosis	11
1.1.5	Tuberculosis treatment	13
1.2	Cytokines play key role in mediating immune response to <i>M. tuberculosis</i>	13
1.3	Tuberculosis and HIV/AIDS co-infection present clinical management challenge	16
1.4	Isoniazid in the treatment and prevention of tuberculosis.....	17
1.5	Host-directed therapies in tuberculosis	18
1.6	Tuberculosis in Tanzania	19
2	AIMS OF THE THESIS	23
2.1	Specific Aims	23
3	MATERIALS AND METHODS	24
3.1	Patient population	24
3.1.1	TB Cohort 1 (Paper I and II).....	24
3.1.2	TB Cohort 2 (Paper III and IV)	24
3.2	Tuberculosis diagnosis and treatment.....	24
3.3	Patient recruitment and follow up.....	25
3.4	Laboratory and clinical procedures.....	25
3.4.1	HIV infection status determination.....	25
3.4.2	Sputum culture for <i>Mycobacteria tuberculosis</i> (<i>M. tb</i>) complex	25
3.4.3	Drug Susceptibility Testing (DST).....	25
3.4.4	Whole Blood Assay (WBA)	25
3.4.5	Cytokine determination by Enzyme Linked Immunosorbent Assay.....	25
3.4.6	Chest radiographs.....	26
3.5	Data processing.....	26
3.6	Ethical considerations.....	26
4	RESULTS AND DISCUSSION.....	29
4.1	Burden of tuberculosis mortality.....	29
4.2	Tuberculosis mortality pattern in relation to anti-retroviral therapy initiation	30
4.3	Resistance to isoniazid and tuberculosis treatment outcomes.....	33

4.4	Anti-EBV or anti-CMV interferon gamma reactivity and tuberculosis treatment outcomes.....	35
4.5	Could cytokines provide a rational biomaker for mortality and lung damage risk stratification	37
5	CONCLUSIONS AND RECOMMENDATIONS	40
6	ACKNOWLEDGEMENTS.....	41
7	REFERENCES.....	45

LIST OF ABBREVIATIONS

AFB	Acid Fast Bacilli
AIDS	Acquired Immune Deficiency Syndrome
APC	Antigen Presenting Cells
ART	Antiretroviral therapy
ATT	Anti-Tuberculosis Therapy
CI	Confidence Interval
COPD	Chronic Obstructive Pulmonary Disease
CT	Computerized Tomography
CTC	Care and Treatment Centers
CXR	Chest X-Ray
DOT	Direct Observed Therapy
DST	Drug Susceptibility Test
E	Ethambutol
ELISA	Enzyme-Linked Immunosorbent Assay
EPTB	Extra Pulmonary Tuberculosis
HIV	Human Immunodeficiency Virus
IFN- γ	Interferon gamma
IGRAS	Interferon Gamma Release Assays
IL-1 β	Interleukin 1 beta
IL-10	Interleukin 10
IL-12	Interleukin 12
IL-2	Interleukin 2
IL-21	Interleukin 21
IL-6	Interleukin 6
IL-6R	Interleukin 6 receptor
INH	Isoniazid
IPT	Isoniazid Preventive Therapy
LJ	Löwenstein–Jensen
LPA	Line Probe Assay
LTBI	Latent tuberculosis infection

<i>M.tb</i>	<i>Mycobacteria tuberculosis</i>
MDR	Multi Drug Resistance
MGIT	Mycobacteria Growth Indicator Tube
MNH	Muhimbili National Hospital
MRI	Magnetic Resonance Imaging
MUHAS	Muhimbili University of Health and Allied Sciences
NAAT	Nucleic Acid Amplification Test
NHP	Non-Human Primates
NNRTI	Non-Nucleoside Reverse Transcriptase Inhibitors
NRTI	Nucleoside Reverse Transcriptase Inhibitors
NTLP	National Tuberculosis and Leprosy Program
OP	Optical Density
PBMC	Peripheral Blood Mononuclear Cells
PCR	Polymerase Chain Reaction
PD1	Program cell Death 1
PET	Positron Emission Tomography
PTB	Pulmonary Tuberculosis
RIF	Rifampicin
SIDA	Swedish International Development Agency
SSA	Sub Saharan Africa
TB	Tuberculosis
Th1	T helper 1
TIM3	T cell Immunoglobulin and Mucin domain 3
TNF α	Tumor Necrosis Factor - alpha
TST	Tuberculin Skin Test
WBA	Whole Blood Assay
WHO	World Health Organization
XDR	Extensive Drug Resistant
YLD	Years Lived with Disability
Z	Pyrazinamide
ZN	Ziehl Neelsen

1 INTRODUCTION

“...She is my only child and a mother of a 5 year old child. Will she survive or die? I need her, for I have nothing to offer to the child....” A voice of a crying woman very early on a Monday morning keeps echoing in my mind as one of the very many cases physicians in Tanzania encounter. This is a vivid image of many years from which I find reason, enthusiasm and strength to carry on with research in infectious diseases - particularly, tuberculosis (TB) and human immunodeficiency virus (HIV) infection. The question posed by the sorrowful mother is a difficult one, which unequivocally demands an answer. At this juncture, we are able to provide patient prognosis largely based on clinical and epidemiological evidence – which is associated with a high level of credibility. Nevertheless, enough patients succumb to TB even in the absence of known predictors of bad outcomes such as HIV co-infection and/or multidrug resistance. In Tanzania, routinely collected TB reports show that 90% of patients newly treated for TB in Tanzania will be cured while about 6% will die during therapy ¹. Every life counts – therefore, there is an urgent need to describe and characterize factors that could provide clues to so many unanswered questions in clinical TB. These answers may also aid in reducing mortality by addressing hitherto unknown factors – so we may one day say to such a grieving mother with great certainty that no one dies of TB or at least, very rarely in our setting! If patients with TB survive, the final outcome may not necessarily be all too positive; 74% of them will develop clinical features of chronic lung disease in the last month of their TB treatment ². Yet, in HIV-infected individuals, a previous episode of TB increases the risk of another TB episode and death ^{3, 4}. Healthcare researchers and practitioners therefore have the obligation to learn how to minimize both the death toll as well as lung damage in patients with TB. In this thesis, we explore factors that could be linked with survival and lung damage in patients with TB, paving the way in search of better tools to improve survival as well as quality of life for patients with TB in Tanzania and internationally.

1.1 TUBERCULOSIS

1.1.1 Etiology and pathogenesis

Tuberculosis (TB) is a deadly disease affecting millions of people, and is caused by the bacterial pathogen *Mycobacterium tuberculosis* (*M.tb*), which was first described by Robert Koch in 1882 ⁵. The infection is transmitted through aerosol droplets from an infected individual by coughing, sneezing, singing or mere talking ⁶⁻⁸. However, it appears that cough is most effective in transmitting the bacteria, not only due to the number of infectious particles expelled, but largely due to the small size of the droplets, which are persistently suspended in air ⁶⁻⁸. Transmission risk is affected by contagiousness of the source (infected person), usually determined by the *M.tb* load in sputum and the proximity between the source and other individuals ⁹. The extent of TB disease in the lung - which in turn increases the frequency of coughing – also greatly affects transmission dynamics, alongside the aforementioned factors ^{8, 9}. In this regard, public health strategies to reducing TB must continue to

address improved living conditions and early case detection and treatment¹⁰. Limiting the extent of pulmonary TB disease by reducing host lung tissue damage using adjunct therapies might also be mutually beneficial to achieve a reduction in lung damage and therefore transmission dynamics.

Inhaled *M.tb* droplets will be transported to the respiratory bronchioles where they will be phagocytosed by alveolar macrophages residing in the lower and middle lung zones¹¹⁻¹³. Effective containment of *M.tb* by the host's immune system establishes a latent infection, termed as latent TB infection (LTBI) as opposed to *M.tb* propagation in the host leading to clinical disease (active TB). Therefore, deficiency in innate immune function¹⁴ and/or adaptive immunity heavily compromises control of *M.tb* infection, fosters infection transmission, and increasing possibilities to succumbing to the disease¹⁵. Acquired immunodeficiency due to HIV co-infection is a major contributor to TB-related deaths, and presents an important clinical setting to study immune cell interactions in TB disease^{16, 17}.

The hallmark of TB pathogenesis lays in the initiation; maturation and maintenance of granuloma as a result of the interplay between innate and adaptive immunity¹⁸. Neutrophils are engaged very early on. Activated by *M.tb* products, neutrophils are responsible for the recruitment of leucocytes and promotion of the inflammatory process. T helper 1 (Th-1) response of the adaptive immunity is central to the formation and maturation of the granuloma¹⁹. The typical TB granuloma, as seen in figure 1, has central caseous necrosis surrounded by different types of macrophages; (epithelioid, multinucleated - Langhans, foamy) all encircled with a layer of lymphocytes^{18, 19}. In non-human primates (NHP) TB granuloma formation has been shown to begin at the hilar lymph nodes after three weeks of infection thereafter involving the regional thoracic lymph nodes and subsequently the lungs approximately four to six weeks following infection²⁰.




Figure 1: A. Granuloma architecture from tissue histological section (Source: Gil et al PlosOne 2010) B. Schematic cellular representation of TB granuloma (Source: Guiradoe et al 2013) both images are taken from reference¹⁸

Disruption of the granuloma results into cavity formation in lungs. Consequently, TB lesions observed is a mixture of inflammatory and tissue healing processes characterized by mixed

immuno-physiological features such as; consolidation, calcification, cavity formation and fibrosis, as well as immune cell infiltrates comprising lymphocytes and myelocytic cells ²⁰. Although TB primarily affects the lungs, *M.tb* can disseminate through the lymphatics to the rest of the body ^{20, 21}.

1.1.2 Global epidemiology of tuberculosis

It is estimated that about 1.7 – 1.9 billion people worldwide harbour LTBI ²²⁻²⁴. The recent global burden of disease estimates that latent tuberculosis is the second most prevalent infection globally, responsible for 1 out of 4 years lived with disability (YLD) in 2016 ²⁴. In general, 10% of individuals with LTBI will develop TB disease at some point in their lifetime ²⁵; the risk is higher in HIV co-infected persons and other in immunosuppressive scenarios such as cancer, transplantation and diabetes ^{26, 27}. Males carry a disproportionately larger burden of TB disease, and are presumptively the source for propagating the infection ^{28, 29}.

According to global statistics, it is estimated that in 2015 there were 10.4 million new patients who developed TB ²³. More than 80% of these cases are in 30 countries classified as high burden TB countries, mainly in Sub-Saharan Africa (including Tanzania) and Eastern Europe, South East Asia and China ²³. The estimated global TB burden is represented on figure 2 ²³.




Figure 2: Estimated global tuberculosis incidence rate 2015 (Source: The WHO - global tuberculosis report 2016)²³

Drug-resistant TB, largely comprising multidrug resistance (MDR-TB) and extensive drug resistance (XDR-TB) in addition to HIV co-infection are the major drivers of the global TB epidemic as well as severe TB disease and increased mortality^{23, 24, 30}. MDR-TB is defined as resistance to at least isoniazid (INH) and rifampicin (RIF)³¹, while XDR-TB is defined as resistance to INH, RIF and any fluoroquinolone i.e. moxifloxacin, levofloxacin, gatifloxacin and at least one of three injectable second-line drugs i.e. kanamycin, capreomycin and amikacin³¹. The global prevalence of MDR-TB is estimated to be 3.3 – 3.9 %^{23, 32}, mainly centred in the South East Asia region as well as China²³.

Re-occurrence of the TB epidemic coincided with the emerging of HIV epidemic in the early -1990s²⁶. Globally, there were approximately 0.5 million HIV-infected individuals with TB reported by the World Health Organisation (WHO) in 2016, while circa 15% of notified TB patients who underwent HIV testing had HIV co-infection²³. As depicted in figure 3, the prevalence of HIV among patients with TB has the highest impact in Sub-Saharan Africa (SSA) where about 80% of TB/HIV cases are usually reported²³. In 2015, the prevalence of TB/HIV co-infection was 36% in the SSA region, particularly high in South Africa^{23, 24}. Only 58% of the estimated MDR-TB cases are treated, while only 55% of all reported TB cases in 2015 had an HIV test with even fewer patients receiving Antiretroviral Therapy (ART)²³. Missed opportunities for treatment such as these should be minimised as we get closer to ending TB by 2030³³.




Figure 3: Estimated HIV prevalence among patients with tuberculosis in 2015 (Source: The WHO - Global tuberculosis report 2016)²³

In 2015, a total of 1.2 million deaths were due to TB; the vast majority had drug-sensitive TB (1.1 Million) while MDR/XDR-TB contributed to a further 0.1 million deaths³⁰. Figure 4 shows the estimated global number of incident TB cases and mortality trends from 2000 to

2015²³. Multiple efforts from different stakeholders have resulted in a reduction of TB cases as well as mortality among tuberculosis patients (figure 4). However, the WHO estimates the trend of decline is inadequate to attain the anticipated goals to end tuberculosis by 2030²³.




Figure 4: Global trends of estimated incident TB cases and mortality 2000 – 2015 (Source: The WHO – Global TB report 2016)²³

1.1.3 Clinical manifestation of tuberculosis

Predominantly a pulmonary disease (pulmonary TB, PTB) in adults, TB has the potential to spread to and affect any part of the body (extra-pulmonary TB, EPTB), and is therefore a systemic disease^{31, 34}. Clinical presentation of patients with TB is variable depending on the affected organ and immune response to the infection. In general, symptoms in PTB include cough, sputum production, low grade fever, night sweats, weight loss, difficulty in breathing and haemoptysis^{35, 36}. These symptoms are non-specific and can be present in many other diseases. On the other hand, it is not uncommon to find disseminated TB presenting with non-classical features or mere asymptomatic among immuno-compromised individuals³⁷. Therefore, in addition to clinical features, accurate and easy-to-use points of care TB diagnostics are highly necessary^{36, 38}.

1.1.4 Diagnosis of tuberculosis

TB diagnosis requires confirmation of the presence of *M.tb*. This can be achieved through bacteriological methods or molecular techniques such as Nucleic Acid Amplification Tests (NAAT)³⁹. Conventional acid fast bacilli (AFB) staining on smear microscopy is a commonly used methods in the clinics in resource limited setting. *M.tb* culture on liquid or solid media is the current gold standard for TB diagnosis³⁹. Molecular methods such as line probe assays (LPA) or Gene Xpert MTB/RIF are instrumental in identifying both TB and drug resistance patterns for clinical decisions³⁹⁻⁴². The Gene Xpert MTB/RIF is an automated, real time Polymerase Chain Reaction (PCR) test that has revolutionized the diagnosis of TB and RIF resistance, thus recommended as standard of care in the context of high TB and HIV^{38, 43}.

Interferon gamma release assays (IGRAs) have been used for diagnosis of LTBI in several countries but are not able to distinguish between active disease and latent infection ⁴⁴. Thus they may not be so useful in countries with high TB burden ⁴⁴. Another challenge with IGRAs is the high variability between tests, making them rather difficult to use for diagnosis and/or monitoring of TB in high burden settings ^{45, 46}.

Radio imaging techniques especially X-ray have been used in the evaluation of patients with TB both to assist in diagnosis, gauging the extent of tissue damage and monitoring treatment progress by assessing tissue healing ³⁴. There are various imaging features of PTB including combinations of consolidation, fibrosis, calcification, cavity, nodulation or fibrosis ⁴⁷. These features are not pathognomonic; there are other disease/pathologies such as sarcoidosis, lung adenocarcinomas and chronic granulomatous disease that mimic TB features on radio-imaging ⁴⁸⁻⁵⁰. However, in settings of smear-negative PTB or TB osteomyelitis, plain radiographs have been useful ^{36, 48}. Unfortunately until recently, there were no simple, validated tools that allowed objective comparison of radiological images between TB patients as well as same patient at different time points during treatment ⁴⁷.

Other imaging methods such as magnetic resonance imaging and computed tomography (CT-scan) with or without positron emission tomography (PET) are essential for the diagnosis of EPTB ⁵¹⁻⁵³. PET-CT and PET Magnetic Resonance Imaging (MRI) are increasingly used in clinical settings especially in TB of the bone and mimics of solid tumours ^{51, 54} but have also been used in research settings as for diagnostic evaluation and/or clinical assessment of cure ^{50, 55}.




Figure 5: Chest x-ray of patient with pulmonary tuberculosis
(Case courtesy of Dr Hani Salam, Radiopaedia.org, rID: 12437)

1.1.5 Tuberculosis treatment

Assessment of anti-TB drug resistance prior to initiating treatment is essential for appropriate management. Treatment of drug-sensitive TB takes approximately six months, and comprises 5mg/kg isoniazid (INH), 10mg/kg rifampicin (RIF), 25mg/kg ethambutol (E) and 15mg/kg pyrazinamide (Z) for 2-3 months followed with 5mg/kg isoniazid and 10mg/kg rifampicin for the remaining time⁵⁶. For re-treatment, patients are given a daily injection of streptomycin (15mg/kg injection) in addition to RIF, INH, Z, and E; for 2-3 months, depending on sputum conversion. It is therefore important to preserve the potency of isoniazid and rifampicin, which form the cornerstone in the treatment of drug-sensitive TB. Although treatment success rate globally is usually above 85% in most cases, there is a need to accelerate the development of new TB drugs to cater for *M.tb* drug resistance strains.

The treatment of MDR-TB is complex, requiring the use of many drugs which are less potent and often accompanied with unpleasant side effects^{42, 57, 58}. The conventional treatment duration for MDR-TB is a minimum of 20 months as guided by *M.tb* culture conversion⁴². Six to seven out of ten patients with MDR-TB will be treated successfully after 20-24 months^{23, 59}. The need for shorter and more effective treatments regimen led to the introduction and inclusion of the new anti-TB drugs Delamanid or Bedaquiline to TB drug regimens⁴², which aim to achieve successful MDR-TB treatment in 9 – 12 months. Both drugs have shown faster and higher culture conversion rates, albeit with associated side effects e.g. arrhythmia. About 80 - 97% of patients with MDR – TB achieve negative cultures, within six months of treatment with Delamanid or Bedaquiline^{60, 61}.

1.2 CYTOKINES PLAY KEY ROLE IN MEDIATING IMMUNE RESPONSE TO *M. tuberculosis*

The success of *M.tb* to establish infection depends on its ability to multiply within the alveolar macrophages and disseminate by counter-acting the host response to control its propagation^{11, 12, 20, 62}. Cytokines are soluble substances that mediate the cross talk between immune cells as well as between non-immune cells, and play an important role in relation to host control of TB through enhanced macrophage as well as lymphocyte activity and granuloma formation^{14, 63}. Many cytokines have been studied in relation to *M.tb* control; this interplay is complex and not completely understood, involving both the innate and adaptive immune cells, as depicted in figure 6^{14, 63, 64}. Several key cytokines have a primary role in *M.tb* pathogenesis and control: interleukin 12 (IL-12), IL-18, interferon gamma (IFN- γ), tumour necrosis factor alpha (TNF- α), IL-1 β , IL-6, IL-17 and IL-2, as observed in animal models of TB and very importantly in patients with TB^{62, 65, 66}.




Figure 6: Interaction of the innate and T helper 1 cytokines following exposure with *Mycobacterium tuberculosis* infection (Source: Cooper Mucosal Immunol, 2011)¹⁴

The interaction between *M.tb* and macrophages induces a series of events leading to production of IL-12 and IL-18⁶⁷⁻⁷⁰. IL-12 is important for the activation of immature dendritic cells as antigen presenting cells (APCs), which will be prompted to translocate from the site of infection to the regional lymph nodes to present *M.tb* antigens to T cells, priming of the adaptive immune response⁶⁶. Priming of CD4+ T helper cell response follows antigen presentation in mediastinal lymph nodes about 10-14 days after the infection^{20, 71}. Cytokines resulting from T helper 1 (Th1) pro-inflammatory responses, IFN- γ , TNF- α , IL-2 and possibly IL-17, in some cellular subsets are crucial in host defence against *M.tb*^{66, 72-74}.

IFN- γ is produced by activated CD4+ and CD8+ T cells and natural killer (NK) cells, and is important to enhance intracellular killing of *M.tb* by macrophages and dendritic cells – which also leads to antigen processing, presentation and augmentation of T-cell activation⁷⁵. Deficiencies in the IFN- γ pathway are associated with failure to control *M.tb* infection, exaggerated tissue necrosis and death^{15, 75, 76}. IFN- γ responses to specific pathogenic *M.tb* antigens has been utilised in the clinical diagnosis of LTBI in low incidence settings. However, its clinical utility as a diagnostic tool for active TB or gauge clinical response remains to be determined^{46, 77}.

During *M.tb* infection, TNF- α is produced initially by monocyte-derived cells, and later by T and NK cells following activation^{65, 78}. Together with IFN- γ , TNF- α potentiates macrophage

phagocytic activity to engulf *M.tb* bacilli and promote apoptosis, which also amplifies the adaptive immune response signals⁷⁹. TNF- α is also important in promoting neutrophil aggregation and is essential for the formation and maintenance of effective granulomas⁶⁵. A fine-tuned balance of TNF- α function is important in modulating TB pathology. Higher levels of TNF- α have been shown to correlate with severe lung pathology^{80, 81}. Alternatively, TNF- α blockade perpetrated increase in *M.tb* load, severe lung necrosis as well as early death in mice⁸¹ and TB reactivation in humans⁸². Interestingly, good clinical responses to anti tuberculosis therapy (ATT) is associated with reduction in TNF- α levels⁷³. In patients treated with anti-TNF- α who eventually develop TB, sudden discontinuation of anti-TNF- α therapy may be associated with severe exacerbation and further lung destruction⁸³. Taken together, TNF- α is an important mediator in TB pathogenesis and its function needs to be well regulated for desirable clinical outcomes.

IL-1 β has also been shown to be important in *M.tb* infection^{14, 81, 84}. The mechanism of IL-1 β -mediated effects in TB pathogenesis is not very clear; however signalling via toll like receptors (TLR) in conjunction with IL-1 β production and activation of the adaptor molecule MyD88⁸⁴ has been shown in TB. IL-1 β is important in recruiting perivascular phagocytes and promoting inflammation and granuloma formation⁸⁵. Mice lacking IL-1 β showed inability to produce IL-12 and TNF; these mice had severe lung disease and died despite a well preserved adaptive immune response⁸¹. Host-directed therapy targeting IL-1 β has been proposed in TB, which reflects the dual nature of this cytokine in maintaining immunological balance in disease⁸⁶.

The importance of IL-17 in TB has been noted in its ability to promote quick trafficking of memory CD4+ T cells from lymph nodes into the lungs for early *M.tb* control⁸⁷. In NHP model of human TB, animals that are able to control TB showed up regulation of IL-17 pathway genes⁸⁸. Similarly, whole blood levels of IL-17 were lower among children with TB compared to healthy controls⁸⁹.

Pleiotropic cytokine such as IL-10 and IL-6 are important in modulating the severity of TB disease but simultaneously allow enough pro-inflammatory activity for successful TB control⁶⁴. Elevated levels of IL-10 is documented in both lung and serum of patients with TB; however its correlation with clinical presentation or outcome is variable⁹⁰. Levels of IL 10 were significantly higher in children with TB disease compared to healthy individuals⁸⁹. It seems plausible that some anti-inflammatory effect of IL-10 is necessary in order to eradicate *M.tb* in the granuloma⁶⁴, since a generally requirement for pro-inflammatory activity exists⁹¹.

IL-6 is a pleiotropic cytokine with both pro- and anti-inflammatory effects that is produced by many cell types including monocytes, macrophages, dendritic cells, mesenchymal stromal cells and fibroblasts^{92, 93}. Patients with active TB have higher levels of IL- 6 compared to healthy controls, and their IL-6 quantities directly correlated with systemic disease, mycobacterial load in sputum and lung injury⁸⁰. IL-6 signals via its receptor, IL-6R in association with gp130, both of which are expressed on T and B cells, hepatocytes and

fibroblasts among others. IL6R-bound IL-6 can also promote trans-signalling, whereby the IL-6-IL6R complex would bind to gp130 on cells to initiate signalling. IL-6 signalling induces the release of several liver proteins, T-cell differentiation, B-cell activation and antibody production, secretion of vascular endothelial growth factor (VEGF), platelet generation and collagen production among others⁹².

Preclinical studies indicate that IL6 may be important for survival during TB disease. Mice deficient for IL-6 succumbed to *M.tb* but not to *M. bovis* BCG challenge³⁰. In one study it was reported that *M.tb* induces IL-6 to inhibit macrophage response to IFN- γ and thereby dampen the adaptive immunity⁹⁴. Blocking IL-6 trans-signalling did not affect protection in *M.tb*-infected mice but is associated with control of inflammation²³. However, complete blocking the IL-6R would reactivate latent TB⁸³. Thus, abrogation of cytokine production from the start cannot be an option but rather modulation of their dynamics during disease. Since *M.tb* infection presents with a wide spectrum of disease (latent vs active; early active disease vs late), cytokine secretion may vary accordingly and studies in mice can't directly be translated to humans. Further clinical investigation among humans with TB is warranted to unfold the role of IL-6, and whether anti-IL-6/IL-6R therapies may be beneficial to reduce lung damage, manage immune reconstitution inflammatory syndrome (IRIS) in patients with or without HIV and optimizing MDR/XDR-TB treatment outcomes^{80, 95}. It is evident that cytokines play key role in TB pathogenesis, but their role of as markers of disease severity and/or clinical response is yet to be precisely elucidated^{46, 77}.

1.3 TUBERCULOSIS AND HIV/AIDS CO-INFECTION PRESENT CLINICAL MANAGEMENT CHALLENGE

The burden of TB among HIV-infected individuals is high; up to 40% of HIV-infected individuals developed TB in the SSA region^{96, 97}. Despite these high TB rates, 30 – 50% of TB cases are diagnosed post mortem⁹⁷⁻¹⁰⁰. HIV-infected individuals are at an increased risk for TB, exhibiting both true relapse and re-infection presenting with mixed *M.tb* strains¹⁰¹. HIV mainly infects CD4+ T cells, dendritic cells and macrophages¹⁰², resulting in impaired control of infections such as *M.tb* due to quantitative and qualitative dysfunction of the macrophage and CD4+ T-cell responses¹⁶. Chetty and colleagues demonstrated that HIV/TB co-infected individuals produce less CD4+ and CD8+ T-cell specific cytokines compared to HIV mono-infected individuals suggesting that the impaired immune function could be the cause for inability to control *M.tb*¹⁰³. On the other hand, impaired control of *M.tb* among HIV-infected individuals, who are antiretroviral therapy (ART) naïve, has been associated with immune cell exhaustion and autophagy^{104, 105}. CD4+ T cells from patients with HIV/TB have been shown to express markers of cell exhaustion (T-cell immunoglobulin and mucin domain 3 (TIM3) and programmed cell death 1 (PD-1)) and exhibited reduced control of *M.tb* infection compared to T cells from healthy controls; this phenomenon was reversed with blockade of TIM3/PD1¹⁰⁴. HIV-infected macrophages demonstrate increased IL-10 production and inhibition of apoptosis¹⁷ which is restored with early ART¹⁶. Overall the

number of incident TB cases has an inverse relationship as the duration of ART increases after initial immune reconstitution phase^{4, 106, 107}.

Diagnosis of TB among HIV-infected individuals presents an enormous challenge to clinicians, often exhibiting atypical symptoms and negative microbiological and imaging results^{37, 108}. The invention of Xpert MTB/RIF has improved TB case detection compared to smear microscopy, with detection rates increased by up to 60% among patients with smear-negative TB^{109, 110}. Yet, about 48% of smear-negative TB/HIV patients will be missed by Xpert MTB/RIF system¹¹⁰. This unmet gap calls for urgent invention of rapid and more sensitive diagnostic protocols.

HIV and TB co-infection also presents challenges during treatment. Treatment of HIV-infected individuals with ART demands careful planning for patients with TB co-infection. Concomitant administration of ATT and ART often results in unwanted drug interactions, increased toxicity and IRIS^{111, 112}. RIF, a key first line anti-TB drug, reduces serum concentration of most non-nucleoside reverse transcriptase inhibitors (NNRTIs), integrase inhibitors as well as protease inhibitors (PIs)¹¹³⁻¹¹⁷. Serum levels of nevirapine and rilpivirine are greatly reduced to sub-therapeutic levels, so the combination of any of these NNRTIs is usually avoided in combined ATT and ART^{114, 117}. Another anti-retroviral drug, efavirenz is affected to a lesser extent than nevirapine, therefore dose adjustment greater than 600mg daily is not usually recommended in general practice¹¹⁸⁻¹²⁰. Rifabutin is a rifamycin of choice for patients with TB/HIV co-infection when PIs are considered¹²¹; however, efavirenz is expensive and not available in low-income countries where TB/HIV burden is high. In contrast, ritonavir-boosted PI decreases rifabutin to sub-therapeutic levels, necessitating dose adjustment to 150mg daily when the combination is prescribed¹²¹. Availability of newer anti-TB drugs such as delamanid provides a broader avenue for choice with fewer drug interactions yet good treatment outcomes although access may be another challenge in resource-limited areas¹²².

Scale-up of ART in the SSA region has tremendously improved survival of patients with or without TB¹²³. However, despite the gains, the risk of TB is highest during the immune reconstitution phase of ART usually the first three to six months of ART and decreases thereafter¹⁰⁶. Multiple reasons therefore contribute to less successful treatment outcomes among TB/HIV co-infected individuals compared to individuals with TB. Therefore, HIV co-infection is often a mandatory component which needs to be accounted for in TB research and clinical practice.

1.4 ISONIAZID IN THE TREATMENT AND PREVENTION OF TUBERCULOSIS

Isoniazid is a potent anti-TB drug with early bactericidal effect^{124, 125}. As a first line drug of choice, INH in combination with RIF is used in standardised regimens to shorten treatment duration of drug-sensitive TB⁵⁶. In addition, INH is recommended as standard of care for TB prophylaxis among recent tuberculin skin test (TST) converters (with or without history of close TB contact), children who are contacts of patients with PTB or HIV infected individuals

after exclusion of active TB disease^{38, 126, 127}. Indeed, INH prevention therapy (IPT) is one of the three pillars of TB prevention in addition to intensified case finding and infection control³³.

The benefit of IPT in preventing mortality and reducing TB occurrence and recurrence among HIV patients is well established in the presence or absence of ART¹²⁸⁻¹³⁰. However, there are some practical uncertainties that need to be sorted out: the ideal duration of IPT for HIV infected individuals, exclusion of active TB among HIV infected individuals in resource-limited settings, sub-optimal compliance to IPT treatment and possibilities of emerging resistance in the long term¹³¹⁻¹³⁴. Compliance to IPT is necessary to prevent mortality¹³⁵. Variable non-compliance rates have been shown, ranging from 2 – 36% for 6 months IPT¹³⁶⁻¹⁴⁰. Worse compliance has been reported among female sex workers, among whom IPT non completion rate was 61%¹⁴¹. These challenges therefore present impediments to the proposed benefits of continuous IPT for HIV-infected individuals¹²⁹, more so when resistance to INH has been linked to non-compliance¹⁴².

INH resistance is the most common type of resistance among anti-TB drugs in clinical use^{32, 143, 144}. Globally, INH resistance does not show a decreasing pattern like that seen among patients with MDR-TB in the United States¹⁴². The median global prevalence of INH resistance is 13% but rates as high as 60% have been reported in some countries^{32, 143}. Mathematical models predict increased INH resistance when community-wide IPT programs are accounted for^{132, 145}.

There are variable treatment combinations to manage INH-resistant TB, in some settings, the treatment of choice is left to the discretion of the clinician based on drug susceptibility testing (DST)^{146, 147}. The WHO recommends using the RIF, Z and E regimen for 6–9 months¹²⁷. Treatment of patients with non-MDR INH resistance shows variable outcomes. Some reports show similar success rates compared to drug-susceptible TB¹⁴⁸⁻¹⁵⁰, while others show less favourable outcomes^{146, 147, 151}. It is therefore essential to monitor INH resistance rates so as to provide evidence for programmatic decision-making for preserving drug efficacy. Notwithstanding high resistance rates for INH, standardised regimens for treating INH mono-resistant TB need better optimisation.

1.5 HOST-DIRECTED THERAPIES IN TUBERCULOSIS

Host immune responses are immensely responsible for disease progression as well as severity of lung damage in patients with TB¹². Modulation of the host response to *M.tb* infection and during TB disease forms the mainstay of host-directed therapies against TB^{152, 153}. These approaches are designed as adjunctive therapies to current anti-TB drug regimens to reduce the extent of tissue damage and improve treatment outcomes¹⁵²⁻¹⁵⁴. Safe and efficacious interventions have been shown for drug-susceptible as well as MDR/XDR-TB^{83, 154}; larger clinical trials for optimization as well as to provide newer options are needed for drug sensitive and drug resistant TB. Therefore, a further understanding of the immune

correlates of severe lung injury and poor outcomes would be important to better understand TB pathogenesis, paving way for better therapies.

1.6 TUBERCULOSIS IN TANZANIA

Tanzania is one of the high burden TB countries according to global statistics^{23, 30}. TB prevalence in Tanzania is estimated to be 249 – 293 per 100 000 adults and the most affected population groups are the elderly, adult males and inhabitants of rural areas^{155, 156}. The distribution of TB prevalence in Tanzania is provided in Figure 7.




Figure 7: Prevalence of TB in Tanzania by cluster (Source: National Tuberculosis and Leprosy Program - The first National tuberculosis prevalence survey in United Republic of Tanzania final report – Ministry of Health and Social Welfare)

Country notification for all forms of TB in Tanzania for the year 2014 was 63,151 ¹. This number translates to a TB case notification of 142 per 100,000 population and 48% case

detection rate compared to the estimated prevalence ^{1, 29}. Increasing age is a strong correlate of TB at population level ²⁹; however, TB cases among the elderly are disproportionately less notified in the health system (Figure 8) ^{1, 155}.

In Tanzania, all patients with TB are provided access to HIV testing services with the possibility to refuse. The HIV screening acceptance rate among TB patients was 88% in 2014. Among those tested for HIV, 19890 (36%) had TB/HIV co-infection and 87% of the identified TB/HIV patients started on ART within the first three months of diagnosis ¹.

The national prevalence of HIV among Tanzanian adults (15 – 49 years) is 5.1%. The proportion of HIV-positivity is slightly higher in urban (7.2%) than rural (4.3%) areas, and among women (6.2%) compared to males (3.8%) ¹⁵⁷. Until 2016, a total of 894,356 people (children and adults) living with HIV (PLHIV) were receiving ART ¹⁵⁸. Due to the increased risk for TB among PLHIV, HIV clinics are important entry points to TB care and vice versa; in Tanzania. TB screening is done using a simple symptom-based tool followed by detailed investigation whenever TB is suspected ^{35, 36}. For TB prevention among PLHIV, IPT is provided at 300 mg dose daily for six months ³⁶.




Figure 8: Age distribution of Tuberculosis notifications in Tanzania (Source: National Tuberculosis and Leprosy Program Annual report for 2014)

Drug-resistant TB is less common in Tanzania, presenting in the range of 0.2 – 1% and 4% for new and retreatment patients respectively ¹⁵⁹. The national health system, however, reported 142 patients with MDR-TB in 2014 ¹. Based on the national survey, about 631 new TB patients presumably have MDR-TB (1% of 63,151 patients with TB notified in 2014) ¹⁵⁹. Therefore, the majority of patients with MDR-TB in Tanzania may die before diagnosis. Increasing the access to rapid molecular diagnostic tools may improve detection of MDR-TB cases in the country.

Diagnosis of TB in Tanzania is mainly achieved by microscopy at primary health centres. Recent reports show that microscopy services are available in 945 facilities across the country¹. Regional referral facilities and several district facilities provided total of 67 Gene Xpert MTB/RIF services in the country in 2015 (compared to 3,500 facilities health facilities treating patients with TB)^{1, 160}. Zonal and national referral laboratories offer culture and molecular technology platforms such as LPA and Gene Xpert MTB/RIF¹⁶⁰.

Tanzania adheres to the WHO treatment guidelines for the clinical management TB^{36, 41, 42}. Treatment is supervised via directly observed therapy (DOT). Patients have two options to choose from: facility DOT, which is observed by healthcare workers at the nearest clinic daily or community DOT, which is supervised by a pre-identified treatment supporter at the patient's residence^{36, 161}. Patients on community DOT have been shown to have an ATT adherence rate of 96%, and in the initial assessment community DOT had similar cure rates but higher treatment success compared to facility DOT¹⁶². The treatment success for smear-positive TB in Tanzania is about 90%, and overall 3,650 (5.6%) patients with of all forms of TB notified in 2013, died at some point during the treatment period¹. Further analysis aimed at improving treatment outcome is of prime importance to save approximately 4000 patients with TB estimated to die per year in Tanzania.

2 AIMS OF THE THESIS

The general aim of this thesis was to understand further how host immune responses and pathogen factors influence treatment outcomes of patients treated for pulmonary tuberculosis in Tanzania. We explored selected host responses (unstimulated and stimulated cytokine dynamics) as well as pathogen factors (HIV infection, antiretroviral therapy and isoniazid resistant *M.tb*) in the hope of contributing knowledge towards reducing mortality and improving quality of life of patients with pulmonary TB in Tanzania. A conceptual framework for the thesis aims is provided in figure 9.

2.1 SPECIFIC AIMS

1. To quantify the mortality burden among patients initiating first line anti tuberculosis therapy in Tanzania (Paper I).
2. To investigate the effect of timing of anti-retroviral initiation in relation to commencement of anti-tuberculosis therapy on the outcome of patients with tuberculosis in Tanzania (Paper I).
3. To compare treatment outcomes among patients with and without isoniazid resistance in the absence of MDR-TB (Paper II).
4. To explore baseline anti-EBV or anti-CMV interferon gamma reactivity as a predictor of death or survival during treatment for tuberculosis (Paper III).
5. To explore the utility of selected serum cytokine levels as biomarkers for disease severity and predictor of poor treatment outcomes for patients with tuberculosis (Paper IV).




Figure 9: Conceptual framework for the stud aims

3 MATERIALS AND METHODS

Detailed descriptions of the methods used for the papers in this thesis are available in the respective papers a brief description is provided and summarised in table 1.

3.1 PATIENT POPULATION

We employed two prospective observational cohorts (cohort 1 and 2) to answer the objectives of the thesis as depicted in Figure 4. Eligibility into both studies included: Recently diagnosed pulmonary tuberculosis at one of the participating research clinics, no prior history of tuberculosis disease or treatment with isoniazid preventive therapy, intention to remain in Dar es Salaam until completion of their TB treatments.

3.1.1 TB Cohort 1 (Paper I and II)

Recruitment of patients for cohort 1 commenced in October 2010 and follow up was completed by December 2011. Patients were 15 years and older and were recruited from one of the participating tuberculosis clinics. We involved the 14 largest TB clinics from Ilala, Temeke and Kinondoni municipals in Dar es Salaam. The health facilities involved were; Amana, Mwanayamala and Temeke Regional referral hospitals; St Monica Modern hospital; Buguruni, Mnazi mmoja, Magomeni, and Sinza Health centres; Mbagala Rangitatu, Mbagala Kizuiani, Mbagala Tambuka reli, Tandale and Ukonga dispensaries; as well as the Infectious Disease Clinic (IDC) of the Muhimbili National Hospital (MNH).

3.1.2 TB Cohort 2 (Paper III and IV)

This cohort included patients aged 18 years and older recruited in Mwanayamala and Amana Hospitals and Mnazi Mmoja Health Centre between October 2013 and April 2015. We excluded patients known to have malignancies, or immunosuppressive therapy or those with end organ failure.

3.2 TUBERCULOSIS DIAGNOSIS AND TREATMENT

Diagnosis of tuberculosis was made by the attending clinicians according to the Tanzanian guidelines for diagnosis and management of tuberculosis.³⁶ Recruited patients were those diagnosed with pulmonary tuberculosis on the basis of sputum smear positive results. Patients provided spot and morning sputum samples and smears were examined for AFB using Ziehl–Neelsen (ZN) technique by trained laboratory technicians at participating health facilities.

All patients were managed in accordance with country guidelines³⁶. Treatment consisted of daily isoniazid 5mg/kg, rifampicin 10 mg/kg, ethambutol at 15 mg/kg and pyrazinamide at 25mg/kg for 2 months (intensive phase) followed with daily 5mg/kg isoniazid and 10 mg/kg Rifampicin for subsequent four months (continuation phase). An additional month of intensive phase ATT was given in case sputum smear conversion was not attained after two

months of ATT. Thereafter, smear non-convertors were managed as having MDR-TB. Throughout duration of TB therapy, DOT was instituted.

3.3 PATIENT RECRUITMENT AND FOLLOW UP

Consenting patients entered into the study upon first prescription of anti-TB drugs. Interviews were performed by attending clinicians. In addition to routine tests, sputum and blood samples were requested at diagnosis. Patients were interviewed and had repeat tests 2 and 5 months after initiation of ATT.

3.4 LABORATORY AND CLINICAL PROCEDURES

3.4.1 HIV infection status determination

HIV infection was determined using DetermineTM HIV-1/2 (Inverness Medical Japan Co. Ltd, Japan) and Uni-GoldTM HIV-1/2 (Trinity Biotech, Wicklow, Ireland) serially. Enzyme Linked Immunosorbent Assay (ELISA) was employed as third test for any discordance.

3.4.2 Sputum culture for *Mycobacteria tuberculosis* (*M. tb*) complex

Detection of *M.tb* was achieved in either the solid egg based Löwenstein–Jensen (LJ) or liquid media system, BACTEC Mycobacteria Growth Indicator Tube (MGIT) – (Beckton-Dickinson) according to manufacturer’s instruction.

3.4.3 Drug Susceptibility Testing (DST)

Drug susceptibility testing was done using the proportion method. *M.tb* H37Rv strain was used as control. The following drug concentrations were used for DST, 0.2mg/l for isoniazid, 40 mg/L for rifampicin, 4 mg/L for streptomycin and 2 mg/L for ethambutol.

3.4.4 Whole Blood Assay (WBA)

Two millilitres of diluted whole blood was added into 96-well sterile culture plates pre-coated with: TB antigens: ESAT-6, Ag85A, Rv0447c, Rv2957, Rv2958c; common viral antigens: EBNA1, CMVpp65, HSV-1, H5N1, H1N1, HIV antigens: HIV env, HIV gag; and controls negative (RPMI) and positive (PHA). The plates were incubated 37°C in a 5% CO₂ incubator for 7 days when supernatant was transferred into non-sterile 96-well plates and frozen at -80 °C for IFN-γ Enzyme Linked Immunosorbent Assay (ELISA) determination later.

3.4.5 Cytokine determination by Enzyme Linked Immunosorbent Assay

Serum concentrations of unstimulated IFN-γ, TNF-α, IL-2, IL-6, IL-10, IL-17A and IL-21; as well as WBA-supernatant concentration of IFN-γ were determined using ELISA kits (Mabtech, Stockholm, Sweden) according to manufacturer’s instructions. Standard curve was

plotted and was then used to determine concentration of the cytokines based on Optical Density (OD) value.

3.4.6 Chest radiographs

Postero-anterior chest radiographs were done at entry and upon treatment completion for a subset of the study population in cohort 2. Radiographs were assigned scores using a validated tool by Ralph A and colleagues⁴⁷. A minimum score of zero was given where there were no radiographic changes while highest score of 100 was assigned when all zones in both lungs were affected homogenously. Presence of cavity regardless of size and number attracted an additional the score of 40. Consequently, the maximum score for any radiograph in the presence of cavities was 140.

Definitions of chest injury:

Severe lung damage at TB diagnosis: Chest x-ray (CXR) score equal to or above 80 (75th percentile) at TB diagnosis.

Severe lung damage at TB treatment completion: CXR score equal to or above score of 50 (75th percentile) at end of ATT.

3.5 DATA PROCESSING

Case record forms were used to capture demographic, clinical and laboratory data which was subsequently transformed to electronic data using Epi info 6 data capture screen. Analysis was either done by either SAS version 9.3 (paper I) or Statistical Program for Social Sciences (SPSS) version 23 (Paper II – IV) as summarised in table 1.

3.6 ETHICAL CONSIDERATIONS

This doctoral project was based on observation study involving human subjects. The project involved; patient interview, taking extra samples, access to the patients' clinical information relating to their treatment as well as outcome.

Ethical approval was obtained for both cohorts 1 and 2 of this thesis. Patients' autonomy, rights and safety were always safeguarded. Eligible patients received information sheets with study details. Patients were guaranteed of appropriate health care even if they did not consent to participate in the study. The patient's relative or another health care worker unrelated to the study assisted patients unable to read and/or write. Parents or guardian provided consent on behalf of minors (patients between 15 – 17 years). Written consent was provided by signing on special consent forms. Patients were interviewed privately; notes were stored in locked cabinets; and the subsequently created electronic database was stored in password protected computers.

Table 1: Summary of materials and methods for studies used in this thesis

Aim (s)	Study Design & Study population(n = sample size)	Main outcome of interest	Main exposure of interest/ Laboratory methods/read- out	Main statistical analysis methods
1/2	Prospective cohort of Tuberculosis patients (n = 1696)	All cause mortality	HIV Anti-retroviral therapy (ART)	Cox-proportional hazards model
3	Prospective cohort of Tuberculosis patients without multi-drug resistance TB (n = 861)	Unfavorable treatment outcome (Death, treatment failure and loss to follow up)	Isoniazid resistance by MGIT and LJ cultures	Logistic regression
4	Prospective cohort of Tuberculosis patients (n = 234)	Death or survival	Anti-gamma interferon level (pg/ml) to EBV from supernatant on whole blood Assay Anti-gamma interferon level (pg/ml) to CMV from supernatant on whole blood Assay	Mann-Whitney U test
5	Prospective cohort of Tuberculosis patients (n = 234)	Death or survival Severe lung injury at baseline (high x-ray score >80) Severe lung damage post treatment (score > 50)	Unstimulated serum cytokines by Sandwich ELISA on serum IFN- γ , TNF- α , IL-6,IL-21, IL-17 , IL-10	Logistic regression General linear model for repeated measure analysis

4 RESULTS AND DISCUSSION

4.1 BURDEN OF TUBERCULOSIS MORTALITY

Paper I: Int J Infect Dis. 2017 Mar; 56:39-44

A total of 1696 out of 1805 patients with tuberculosis were included in this paper, among them 58 (3.4%) deaths occurred within six months of ATT. We excluded 109 patients because they had missing information such as HIV, outcome and age. The excluded patients were more likely to be males and had history of illicit drugs use; factors linked with increased mortality risk among patients on ATT ¹⁶³. Due to this potential bias therefore, we might have underestimated mortality. In the worst case scenario, if all 109 excluded patients are considered dead, then, mortality would be 9.3%. Thus, it is very likely that the true burden of mortality among patients with TB in our study would be between 3.4 – 9.3%.

The mortality risk was higher among TB/HIV co-infected individuals (41/514; 8.0%) as compared with TB mono-infected patients (17/1182; 1.4%), a phenomenon reported among patients with HIV especially in advanced disease ^{96, 164, 165}. Most probably as a consequence of multiple factors; related to immune suppression, ¹⁶ immune reconstitution or, not mutually exclusive, competing drug toxicity ^{111, 166}.

The median time to death in our cohort was 46 days and at the end of intensive phase of ATT two thirds of all the deaths had occurred. This accelerated initial mortality warrants further investigation in relation to exaggerated inflammation. The initial phase of ATT is associated with exponential killing of *M.tb* resulting in a surge of *M.tb* antigens accessible to the immune system ⁹. Worsening of clinical features following effective ATT; also known as *paradoxical TB*, has been described among HIV infected and uninfected individuals and might be related to exaggerated inflammatory response ^{166, 167}. On the other hand, early deaths may signify advanced disease owing TB diagnostic challenges ³⁷. It could be hypothesized that, late presentation is associated with large TB antigen burden or a mere host exaggerated inflammatory response but this remains unanswered by this study and calls for further studies.

4.2 TUBERCULOSIS MORTALITY PATTERN IN RELATION TO ANTI-RETROVIRAL THERAPY INITIATION

Paper I: Int J Infect Dis. 2017 Mar; 56:39-44

It was apparent that TB/HIV co-infected patients (n = 514) had higher mortality risk thus warranting a much deeper scrutiny in relation to HIV treatment. As shown on table 2; patients who initiated ART 90 days before starting ATT or within first 14 days of ATT had highest mortality (10%), whereas lowest mortality was observed among TB/HIV who had initiated ART after 14 days of ATT (5.0%) (Table 2).

Table 2: Crude mortality pattern in relation to HIV infection and anti-retroviral therapy (Nagu, IJID 2017)

	No. of deaths	No. at risk	Crude mortality rate (%)	Median days to death (IQR)
TB+/HIV	17	1182	1.44	46 (32, 62)
TB+/HIV+; ART >90 days prior to ATT	5	49	10.20	79 (33, 96)
TB+/HIV+; ART ≤ 90 days prior to ATT or ≤14 days after ATT	4	39	10.26	56 (27.5, 72)
TB+/HIV+; ART >14 days after ATT	7	141	4.96	52 (45, 89)
TB+/HIV+; No ART	25	285	8.77	37 (26, 59)
Total	58	1696	3.42	46 (30, 72)

IQR = Inter Quartile Range, TB = Tuberculosis, ATT = Anti Tuberculosis Therapy

In this cohort, less than half of TB/HIV patients (229/514) were initiated on ART during the follow up period. According to Tanzanian guidelines at the time, all HIV co-infected TB patients had to be initiated on ART, preferably after the second week of ATT or within the first two weeks in case of severe immunodeficiency (CD4+ T lymphocytes < 50 cells/μL) ³⁶. The low proportion of HIV infected patients on ART is probably a reflection of a cascade of events including late HIV diagnosis, patients' refusing to take ART, poor referral and lack of integration of HIV and TB services at the time. Integration of HIV and TB services has lately improved ART uptake to 87% but mortality remains at 6% ¹. Since, TB clinics serve as an important entry point to HIV care and treatment, attempts to harness integrated TB and HIV services for appropriate and timely interventions would greatly improve treatment outcomes.

Using multivariate analysis we compared mortality in relation to ART and ATT among TB/HIV co-infected patients, while TB patients without HIV co-infection was the reference group. Mortality risk was lowest (three fold) when ART was initiated more than 14 days after ATT and highest when ART was initiated before or within the first 14 of ATT. Persistent high mortality among TB/HIV patients despite good HIV virological control is yet to be thoroughly understood among patients on ART. Apparently there is heterogeneity in regaining overall immune function after ART initiation despite viral suppression which may be associated with lack of TB control and/or mortality ¹⁶⁵. It suffices to say at this point that,

ART and ATT initiation by themselves are insufficient to avert some deaths among TB/HIV patients on ART. It is thus of interest to further investigate the functional immune reconstitution in relation to clinical recovery and/or possibility to intervene with host directed therapies to reduce these unaccounted mortalities.

Patients with advanced HIV particularly those with TB/HIV need vigilant follow up of clinical, virological and immunological parameters; and prompt action should be taken when required¹⁶⁵. TB/HIV patients, who had initiated ART more than three months prior to ATT in our cohort, had low median CD4+ T lymphocyte count (263 cells/ μ L) and median BMI of 18.8kg/m². These clinical parameters are associated with opportunistic diseases and thus may be synonymous with a failing ART regimen. We did not quantify HIV viremia, neither did we study ART resistance nor the functional CD4+ T cell immune recovery in relation patients who died. This might be regarded as an area to improve the quality our study in order to add further knowledge for even better TB/HIV management. We discuss inflammation in relation to mortality in papers III and IV.

Given the overall high initial mortality, it is plausible to hypothesize that the individuals who died before ART was initiated had been sicker and therefore incomparable to those in the ART group. This hypothesis may not likely in our cohort as patients had similar median CD4+T cell count and the body mass index (BMI) as seen in Table 3

Table 3: Distribution of median CD4+ T lymphocyte counts and body mass index by anti-retroviral therapy among TB/HIV co-infected patients (BMI = Body mass index; IQR = interquartile range) (Nagu, IJID 2017)

	Median CD4+ (IQR)	Median BMI kg/m ²
TB+/HIV+; ART > 90 days prior to ATT	263 (181, 422)	18.8 (17.0 -21.5)
TB+/HIV+; ART \leq 90 days prior to ATT or \leq 14 days after ATT	168 (79, 221)	17.4 (15.4, 19.6)
TB+/HIV+; ART >14 days after ATT	151 (69, 220)	18.8 (16.9, 21.2)
TB+/HIV+; No ART	331 (158, 471)	18.1 (16.5, 20.4)

IQR = Inter Quartile Range, TB = Tuberculosis, ATT = Anti Tuberculosis Therapy


There were yet HIV infected patients who developed TB following ART in our cohort 1 (Table 2). Unmasking TB is a term coined to TB occurring shortly after initiation of ART usually in the first three months of ART^{107, 168}. The increased rates of TB soon after ART are probably a result of TB missed during clinical evaluation prior to initiation of ART a fact that underscores need for intensified TB screening including use of culture and gene Xpert MTB/RIF pre-ART¹⁰⁸. Unfolding TB might explain the occurrence of TB among the patients who had initiated ART around 90 days prior to or immediately after ATT (within 14 days) (Table 2).

The cohort was almost entirely composed of young to middle aged patients, (91% were ≤ 50 years; and two thirds were males (67%), a finding that is in keeping with many parts of developing countries with high burden of tuberculosis ²³. Two important things to consider include i) increased propensity of such patients to infect their family members as well as colleagues at work places even before diagnosis, as determined by mycobacterial load in the sputum ⁹. and ii) reduced productivity due to prolonged illness related sick leave leading to poverty and malnutrition, a vicious circle of tuberculosis. Indeed, poverty and tuberculosis are cousins who must be addressed hand in hand to break the circle ¹⁰. It is through improved living conditions that TB was significantly reduced in high income countries.

4.3 RESISTANCE TO ISONIAZID AND TUBERCULOSIS TREATMENT OUTCOMES

Paper II: J Antimicrob Chemother. 2017 Mar 1;72 (3):876-881

This study included 861 TB-culture positive patients who had both TB treatment outcome and anti-TB drug susceptibility reported, without dual resistance to isoniazid and rifampicin (MDR) as shown in figure 10.



^aMOTT- Mycobacteria Other Than Tuberculosis

Figure 10: Patient flow chart (Nagu, JAC 2016)

Isoniazid resistance was present in among 23 (2.7%) patients, in combinations with streptomycin 2/23(8.7%) or ethambutol resistance 2/23 (8.7%). Patients with rifampicin/isoniazid co-resistance were excluded prior. During six months of follow up, 25 (2.9%) patients died, 10 (1.2%) patients failed treatment and 29 (3.4%) patients were lost from follow-up at the clinics.

A total of 18/23 (78.3%) patients with resistance to isoniazid had successful treatment outcomes compared to 779/838 (93.0%) patients with preserved isoniazid susceptibility.

At multivariate analysis (Table 4) patients with isoniazid resistance isolates were six times more likely to have poor treatment outcome (RR 6.0; 95%CI 1.9 – 18.7; $P < 0.01$). In addition, HIV infection with (RR 2.3; 95%CI 1.0 – 5.2; $P = 0.05$) or without ART (RR 3.1; 95% CI 1.5 – 6.2; $P < 0.01$) was associated with increased risk for unfavorable treatment outcomes. (Table4)

Table 4: Factors associated with unfavorable treatment outcomes among tuberculosis patients initiating first line treatment in Tanzania (Nagu, JAC 2016)

	Univariate			Multivariate ^a				
	RR	95% C.I.		P	RR	95% C.I.		P
		Lower	Upper			Lower	Upper	
Age (years)	1.0	1.0	1.0	0.83	1.0	1.0	1.0	0.85
Male sex	1.4	0.8	2.4	0.27	1.6	0.8	3.3	0.16
Isoniazid resistance	3.7	1.3	10.2	0.01	6.0	1.9	18.7	<0.01
HIV status				<0.01				<0.01
HIV+/ART+	2.0	1.0	4.1		2.3	1.0	5.2	
HIV+/ART-	2.8	1.5	5.1		3.1	1.5	6.2	
Primary school or below	1.8	0.8	3.6	0.13	1.2	0.6	2.6	0.60
Smoking Status								
Never smokers				0.10				0.78
Past smokers	1.8	1.0	3.1		0.9	0.4	2.0	
Current smokers	0.6	0.2	2.7		0.6	0.1	2.6	
BMI (kg/m ²)	0.9	0.9	1.0	0.19	1.0	0.9	1.0	0.32

ART = Antiretroviral Therapy, BMI = Body Mass Index RR = Relative Risk CI = Confidence Interval)

Initial primary isoniazid resistance although has relatively low prevalence, is associated with poor outcome. These patients harboring resistant *M.tb* would otherwise go unnoticed or discovered late during treatment since anti-TB drug susceptibility testing is reserved to those failing first regimen or patients with a relapsed TB in Tanzania. Phenotypic assessment of drug resistance is resource and technically demanding. In many setting Gene Xpert MTB/RIF has simplified and improved detection of both TB and rifampicin resistance^{109, 110}. However, given the fact that resistance to isoniazid far more common than rifampicin^{143, 159}, it is likely that some cases will be missed by Gene Xpert MTB/RIF system. Patients with undetected INH resistant isolates are likely progress to MDR and/or transmit the resistant strain to close contact, consequently defeating one of the three strong pillars of the END TB strategy^{10, 33}. Accordingly, effective mechanisms to identify and appropriate manage patients with INH resistant *M.tb* are warranted. In addition, simple treatment regimens for INH resistant TB ought to be standardized to enable primary medical personnel to easily administer as it is with management of drug sensitive and MDR TB^{42, 56}.

4.4 ANTI-EBV OR ANTI-CMV INTERFERON GAMMA REACTIVITY AND TUBERCULOSIS TREATMENT OUTCOMES

Paper III: Int J Infect Dis. 2017 Mar; 56:136-139

We recruited 274 patients in this cohort; forty (40) patients were excluded in this report as shown in figure 11 below.




Figure 11: Patients flow chart

A total of 21 deaths occurred during the course of follow up while 213 patients were successfully treated. As in cohort 1, we noted in this cohort as well, an accelerated initial mortality since 72% of the patients died within the first three months of ATT.

Patients with tuberculosis who died, exhibited significantly lower median IFN- γ responses to ESAT-6 (0.043), CMV pp65 ($p = 0.035$) and EBNA1 ($p = 0.006$) in the whole blood assay at the time of diagnosis (Figure 12). There was no statistically significant difference in IFN- γ responses against the following antigens: Rv2958c, Rv0447c, PPD, Ag85A, herpes simplex virus 1 (HSV-1) glycoprotein and HIV env. (Figure 12)

The higher IFN- γ responses to ESAT-6 among survivors of TB, is a reflection of good *M.tb* antigen specific T cell response against *M.tb*. Coupled with similar pattern specific T cell responses against tuberculosis (ESAT-6), responses against EBV and CMV antigens may signal a correlation with good immune response against TB. Whether it is a general reflection of a good immune system or rather that a prior sensitization with EBV and CMV antigens prepared the immune system to be more alert for the mycobacterial antigen is yet to be ascertained. It has been reported that non-specific sensitization of alveolar macrophages was an important factor for their activation to achieve better control of TB infection ¹⁶⁹. Prospective experimental models with CMV or EBV prior to TB infection could help decipher this theory. CMV infections has been shown to drive TB immune response towards memory and terminally differentiated effector T cells ¹⁷⁰, cells which play an important role in controlling TB. EBV also induces potent Th1 responses in EBV-positive individuals. Demonstrating CMV/EBV DNA in these patients would be helpful in

interpreting these results in consideration of prior or acute infection which would differ in their immune activation states^{15, 170}.




Figure 12: IFN- γ responses to viral and mycobacterial antigens in whole blood assay at the time of TB diagnosis in relation to survival during treatment with anti tuberculosis therapy (Nagu, IJID 2017)

4.5 COULD CYTOKINES PROVIDE A RATIONAL BIOMAKER FOR MORTALITY AND LUNG DAMAGE RISK STRATIFICATION

Paper IV: Manuscript

At the time of TB diagnosis we found in serum from 234 patients with pulmonary tuberculosis significantly increased amounts of IL6 (median: 3863 pg/ml) compared to the amounts in serum from seven healthy controls (11pg/ml). In addition IFN- γ , TNF- α and IL-10 levels were also upregulated in patients with TB compared to the healthy controls but the difference were not as profound as it were with IL-6. Patients who died had significantly lower median serum levels of IL-6 at diagnosis compared to those who survived and were cured.

The median CXR score at TB diagnosis was 60. Compared to TB survivors, patients who died had lower median CXR score at diagnosis. The CXR scores tended to decrease at the end of treatment among TB survivors; and in addition we observed a significant linear correlation between diagnosis and end of treatment CXR scores ($r = 0.44$, $P < 0.0001$). In summary, patients with more lung damage at diagnosis tended to have more residual damage at the end of treatment. As much as this seems intuitive, the x ray scores could be utilized in stratifying those who have high risk of severe lung damage for an intervention that might reduce this eventuality. Adjuvant treatment with cytokines and particularly IL6 (or anti-IL-6, dependent on the timing of treatment), is proposed as one possible clinically relevant area for further research as discussed later in this chapter.

Increasing serum IL-6 and IFN- γ concentrations were significantly associated with reduced mortality risk. IL-6 levels observed at diagnosis may protect against death, patients who survived TB had significantly increased lung damage at the end of ATT. This observation is in line with previous findings in sera of patients with TB from Bangladesh¹⁷¹ and South Africa¹⁷².

As previously reported, IL-6 is important in clearing large virulent mycobacterial load through IFN- γ induction. IL-6-deficient mice were shown to produce significantly lower IFN- γ and elevated IL-4 levels in response to *M. tb* challenge, and succumbed to infection within 50 days. Conversely, IL-6-deficient mice infected with the *M. bovis* BCG vaccine strain were able to control the bacterial burden almost as efficiently as wild type mice¹⁷³ which may probably signify the importance of IL-6 in *M.tb* but not *M.bovis* BCG strains.

At TB diagnosis, high levels of circulating IL-6 associates with extensive lung tissue destruction among patients with TB, based on CXR findings. This is in keeping with previous findings in patients with pulmonary TB: i) high levels of circulating IL-6 in TB patients with or without chronic obstructive pulmonary disease (COPD), in conjunction with impaired pulmonary function (marked by reduced FEV1 values)¹⁷⁴, and ii) increased IL-6 levels in bronchoalveolar lavage fluid, with extensive computed tomography findings in the lungs¹⁷⁵.

Additionally, the IL-6 levels in serum/plasma at the end of the treatment are not much lower than after two months of ATT. In support of our observations, several clinical studies have shown that IL-6 is upregulated in patients with TB undergoing therapy¹⁷⁶⁻¹⁷⁸. Thus, it appears that the continuation phase of ATT (last four months, daily INH and RIF only), while contributing to microbiological cure, does not promote pulmonary tissue healing in patients with TB who sustained extensive lung pathology, nor does it help reduce the high IL-6 levels present before and during ATT.

Inflammation-induced tissue damage during ATT leads to a decline in life quality in the long term, although the patient effectively clears *M.tb* infection of the lungs^{2, 179}. The link between pulmonary TB and impaired lung function despite completion of therapy has also been reported for patients in South Africa with drug-sensitive¹⁸⁰ or multidrug-resistant TB¹⁸¹. As a consequence, these individuals are denied of a normal lifestyle thereafter, which affects their contribution to the workforce and directly affecting the country's socioeconomic development. This scenario calls for host-directed therapy (HDT) targeting overt inflammation in addition to conventional antibiotics against *M.tb* not only to prevent early mortality but also to improve life quality of individuals surviving pulmonary TB.

Anti-TNF- α therapy with adalimumab (given adjunctively to standard ATT) during severe pulmonary TB has been shown to be life-saving in severe pulmonary TB⁸³. The findings in the current study lay the foundation for future studies of HDT targeting exaggerated levels of circulating IL-6 in pulmonary TB. Monoclonal antibodies targeting IL-6 (siltuximab) and its receptor (tocilizumab) are in the clinic, approved for treating arthritis and Castleman's disease¹⁸². Further clinical trials are underway to test their therapeutic efficacy in other indications i.e. diabetes mellitus, pancreatic, lung, ovarian cancers, myeloma^{183, 184}. One preclinical study in a mouse model of TB showed that blocking IL-6R with a monoclonal antibody allows for reduced lung inflammation, improved control of lung *M. tb* burden and prolonged survival of the animals¹⁸⁵. However, in the preclinical study, the animals were given the anti-IL-6R monoclonal antibody prior to *M. tb* challenge. Since IL-6 production by human monocytes is crucial for the early control of *M. tb* replication¹⁸⁶, and is required for initiation of adaptive immune responses to *M. tb* antigens¹⁸⁷, we propose that HDT directed at reducing circulating levels of IL-6 in patients with pulmonary TB would be best administered after the first two months of intensive TB treatment. The efficacy of this intervention is best studied in line with reduced inflammatory markers in serum i.e. unbound IL-6, C-reactive protein (CRP), tissue scarring (CXR scores) and restoration of lung parenchyma and pulmonary function (spirometry).

PTB is mainly local disease whose pathogenesis is complex and resulting from many factors including innate and adaptive immunity cells, cytokine and chemokines. Picture we deduce from blood therefore, is but a small part of the whole network responsible for TB

pathogenesis and control. Yet a very important suggestion that we deduce is that intervening with cytokine adjuvant therapy in addition to ATT may modulate this network towards better TB control in reducing local inflammation in the lung.

5 CONCLUSIONS AND RECOMMENDATIONS

In Tanzania, mortality among newly diagnosed TB patients during the course of treatment with first-line regimen is between 3.4 – 9.3%. This mortality is higher among patients with TB/HIV and was lowest when ART was instituted after 14 days of ATT. These findings have a great use in real life clinical practice in Tanzania and other similar setting of high rates of TB and HIV co-infections. Tuberculosis clinics have shown to be a major entry point for HIV care since the majority of our patients were first time diagnosed at TB clinics. In current view of “HIV test and treat strategy”, where everyone found to be infected with HIV will be treated promptly, it is prudent to wait until the third week of ATT. For patients who have been on ART before TB was diagnosed, vigilant monitoring of clinical response and timely intervention is essential.

Background isoniazid resistance is not uncommon and has proved to cause unfavorable treatment outcomes in at least 21% of patients with resistance. Current available *M.tb* culture and DST techniques are demanding in terms of infrastructure and skills for country like Tanzania. Research for point of care, technology friendly machines for early detection of isoniazid and/or rifampicin resistance, need to be prioritized and accessed in routine TB care in order to improve treatment outcomes.

Cytokine responses are essential to successful TB treatment as well as reduction of severe lung inflammation. Whereas serum level of IL-6 are comparatively higher than of other cytokines in sera of patients with pulmonary tuberculosis and are predictive of survival; they could also be a cause of severe pulmonary damage in these patients. Adjunct therapy with IL-6 may serve as possible target in reducing lung damage following TB disease. Timing is of essence: and it may very well be that anti-IL-6 treatment (during the consolidation phase of the TB disease, associated with lung damage), could be a clinically viable option; similarly to the ‘double-edged sword’ described for anti-TNF alpha blockage during different phases of TB disease development⁸³

More studies on inflammation and lung damage, including linking them to lung function tests; longer term cytokine follow for chronic lung inflammation including (COPD); Trials of IL 6 for TB patients as adjuvant therapy especially during the continuation phase of ATT are really essential. Further gauging anti-CMV/EBV responses in *M.tb* disease in patients with known CMV/EBV viremia would certainly add value.

6 ACKNOWLEDGEMENTS

Indeed it has been a long journey, full of adventures and certainly educative in many ways. I will always be grateful to be part of Karolinska Institutet as a doctoral student. Many people have contributed to making this thesis, I would like to thank you all.

To my supervisors;

Professor Markus Maeurer my main supervisor, thank you so much for accepting me as your PhD student at TIM. I am sure it wasn't so easy for you to fit my short stays in Sweden with your busy schedule. You accorded me with the necessary tools and support as much as possible. Mark, you are a talented scientist with unwavering dedication to excellence and you are a gifted teacher. It was a privilege to be your student and hopefully you shall continue to mentor me remotely.

Professor Ferdinand Mugusi, you are an incredible supervisor, passionate clinician and brilliant researcher. You played the double role of a supervisor and a mentor; I am grateful for your untiring support and numerous opportunities to learn. You allowed me to venture into the unknown to test hypotheses; and never ceased to guide. It was an honor to be your PhD student, and even more exciting that I will continue to shadow you in respiratory and infectious disease clinical practice.

Professor Matee, you have known me a long way in research training. You were my supervisor for master's thesis as well as this Ph.D. training. Thank you for all the support in setting up the laboratory and immense help in reviewing my manuscripts.

Dr Rebecca Axelsson, I am grateful for all the help with the immunology experiments. Distance between Tanzania and Sweden was reduced to minimum because you were just a skype call away to make things work out. Thanks Rebecca for taking part of your maternity leave to help with paper work and in reading cover story of this thesis.

I would also like to thank former supervisors Dr Isabelle Magalhaes and Antony Rothfuchs for accepting to be co-supervisors and checking on me. Isabelle, thank you for assisting with the project start-up; and for the admission seminar rehearsal.

To my Mentors;

Professor Muhammad Bakari, you have been a key person to my career development and PhD training. You associated me with SIDA project and never hesitated to intervene when things did not seem to work. Thank you for the initial research skills as my master's thesis main supervisor, without which the beginning of my Ph.D. would have been harder.

Professor Wafaie Fawzi, thank you for being my mentor for the Global Health Scholar program which provided major part of the funds to carry out research for the first two papers in this thesis. I appreciate your support and encouragement as well as the many learning opportunities I obtained through you.

Sir Alimuddin Zumla, it has been a great privilege to know, meet and work with you. Thank you for generously sharing your vast knowledge and experience in tuberculosis with me. I appreciate your help with manuscript revisions, your advices and support.

Professor Pallangyo, you read my manuscripts and provided so many valuable inputs. Thank you for all the advice, encouragement and support during this training.

Dr Johnson Lwakatare, being the next door office neighbor to my home supervisor, it was convenient for me but I am sure not for you. You accommodated my unscheduled visits and offered advice. Thank you.

Professor Julie Makani, you are amazing in so many ways; brilliant researcher, dedicated clinician and great teacher. You have been very inspirational and helpful in my training, thanks Julie.

Dr Martin Rao, you are a great person, thank you for who you are and for all the help you provided me with. Your knowledge in TB as well as scientific writing skills are remarkable; and have been very useful during my training. You were a key contact here in Stockholm while I was in Dar es Salaam and a very good link with Mark. Thanks for helping with receiving and storage of my samples. Thank you for the help in manuscripts and on the thesis as well.

To SIDA, I would like to sincerely thank the government of Sweden through SIDA who funded my training. It was a great opportunity to learn and network.

Special thanks to **Professor Gunnel Biberfeld** and **Professor Anna Mia Ekström** HIV-subprogram coordinators for all practical help. Anna Mia, in addition to being SIDA coordinator, you offered advice whenever I called on you. Professor Said Aboud, you were not only SIDA coordinator in Tanzania but also a great help in laboratory work.

To Muhimbili University of Health and Allied Sciences (MUHAS); many thanks for your support as an employer. I appreciate your initiative to pursue SIDA training grant. I hope attained knowledge and skills will benefit the university and Tanzanians.

To the patients, you have contributed in the search of knowledge for a good cause. Your sacrifice is honored and I remain indebted to you and all patients. To the patients who died during the study, may you Rest in Peace.

To the department of Laboratory medicine; Special thanks to **Professor Matti Sällberg**, **Professor Antony Wright**, **Professor Moustapha Hassan**, **Dr Jenny Degerholm Langsmo**, and **Arja Kramsu** for your great support throughout training but more specifically, during the final months of my training. I am grateful for all you did to make my defense go through as planned. I am proud to be a trainee in this department.

To the department of Internal Medicine MUHAS; many thanks to the former and current heads of department, Professor Janet Lutale and Dr Paschal Ruggajo and all

colleagues for your advice and well wishes. Special thanks to Dr Grace Shayo, Dr Ewaldo Komba, Professor Eden Maro for taking my responsibilities whenever I travelled.

To TIM members – F73-79 corridor – thank you for the happy moments, I enjoyed the fikkas. Special thanks to **Associate Professor Helen Kaibe**, you efficiently respond to matters with great speed and highest level of precision. Thank you for trouble shooting with my defense application process. **Gunbritt Lindholm**, thank you for fixing access cards and all practical help in the hospital.

To Professor Markus Maeurer's research group members former and current; Lalit Rane, Aditya Lambati, Nancy Alvarez-Corrales, Giovanni Ferrara, Thomas Poirer, Luo (a.k.a. Petter), Liu Zhenjiang, Qingda Meng, Annurupa Nagchowdhury, Esther Schoutrop, Geogia Paraschoudi and Chaniya Leepiyasakulchai – I am grateful to you all. You were wonderful in many ways; from practical issues when I stayed in Stockholm to great scientific discussions. I learnt a lot from our interactions. Thomas, thank you for bearing with my clumsiness in the lab at the beginning; thanks too for helping to secure storage for my samples. Marlene and Moa, I am grateful for all your support with administrative issues.

Co-authors; Dr Lulu Fundikira and Dr Zuhura Nkumbih, I appreciate your help in interpretation scoring chest radiographs. Ramadhani Mwiru –thank you for great help with stastical analysis.

National Tuberculosis and Leprosy Program (NTLP). I would like to appreciate the generous support from the program through the former (Dr Said Egwaga) and current (Dr Beatrice Mutayoba) program managers, which enabled smooth navigation through TB clinics. Dr Egwaga thank you for the Tuberculin PPD reagents.

Research staff: I would like to thank all staff who assisted with the research; all the clinicians and nurses at the study sites; Sr Magreth Matekele and Sr Happiness for coordinating the clinical studies; laboratory technicians: Colman Mchau, Salome Kihampa and Betty Mchaki, Zacharia Mtulo, Edward Shogolo, Bryceson Malleo Daphne Mtunga and Basra Doulla. Special thanks to Gladness Kiwelu and Thekla Mtomoni for assisting with data entry; and to Halfan, for transportation of samples.

SIDA PhD colleagues; Dr Matilda Ngarina, Dr Patricia Munseri, Dr Edith Tarimo, Dr Abel Makubi, Dr Agricola Joachim, Dr Frank Msafiri, Dr Majigo Mtebe, Dr Theodora Daniel, Roselyne Urio, Masunga Iseselo, Dr Goodluck Lyatuu and Dr Fadhluni Alwy; thank you all for your help. Dr Helga Naburi and Dr Dawa Mushi you were a great friends and good company in Sweden.

Friends in Sweden Associate Professor Anna-Berit Arvidson, thank you for comments on my “kappa”, Rashmi, Lien, Dr Shweta, Dr Muslima (Say hi to Ajazi) and Dr Uzmali I will treasure the memories.

To my Christian community of our Lady of perpetual help – Dar es Salaam, my neighbors; Martha Kamuhabwa and Domitila Makubi thank you for all you have been.

To my friends and family; thank you for your well wishes, encouragement, support and love all of which sustained me throughout the training period. Mamdogo Yacinta, you are a unique person to me; friend, sister and auntie. Thank you for all you did all these seven years, so my stay away from home was not so stressful. Ephrasia, you had to sacrifice a lot to stay home with Jo-Marie and Davidson. Thank you Efuu, it meant a lot. My cousins Esther, Basila, Anna, Patricia and Jullie; thank you for your support. Redimna and Lucy, you made sure kids were as comfortable as possible. Leticia, I left you home almost all the time, your help and leadership are greatly appreciated. Betty thanks for the encouraging conversations over the phone.

To my in laws, you are an adorable family. Emma and Nelson; Janet and Immanuel, Laura and Richard; and Victor, thank you for your understanding and support in many ways. All of you were a phone call away; dada Emma thank you for sending references whenever I called on you. Janet and Laura thank you for making it to school occasions. My mother in-law, aka “fogo” thank you for your encouragement, support and reassuring words which ended unnecessary worries.

To my parents Mary and Joseph Nagu, I will always be grateful to your ever loving parenting. Mama, you believed in me and gave courage and to go on. Thanks for spoiling wajukuu, every bit is appreciated mama and I love you so much. Daddy, you encouraged and supported me all along. It’s a pity you missed it by a few months, all I have is to imagine your smile as it comes to the end. My siblings Neema and Deogratius, we not only share parents but also very special bond. Thank you for being wonderful persons I could count on. Fr. Wolfgang Pissa O.F.M.Cap; a brother and spiritual advisor, thank you for helping to me to conquer my fears and remain calm.

To my family; **Jo-Marie, Davidson and Lawrence**, it couldn’t be easier without you. Sweetheart Jojolito when I started this journey you were only 13 months, you endured lengths of times without mommy but became a responsible and courageous girl. Thank you my daughter for hanging in there.

Davidson, mommy’s boy – Ooh what an encouraging personality? Thank you for all your love and prayers. I still recall the warmth of your hug as you said God will take care of you mama, when I took my first flight to Sweden, thank you Davidson.

Lawrence my husband, you are the one who shouldered everything at the expense of your own professional development – struggling with loneliness, taking care of our children and my frustrations; you saw it all. I had peace to carry out my training knowing you were there. I have no better definition of a gentleman than I find in you. Thank you so much.

To God be the glory, great things he hath done for me throughout this training, may the service I shall provide be accepted as a sign of my gratitude to the Most High.

7 REFERENCES

1. National Tuberculosis and Leprosy Program. National Tuberculosis and leprosy Program Annual report for 2014. Dar es Salaam; 2015.
2. Manji M, Shayo G, Mamuya S, Mpembeni R, Jusabani A, Mugusi F. Lung functions among patients with pulmonary tuberculosis in Dar es Salaam - a cross-sectional study. *BMC pulmonary medicine*. 2016; **16**(1): 58.
3. Gupta A, Nadkarni G, Yang WT, Chandrasekhar A, Gupte N, Bisson GP, et al. Early mortality in adults initiating antiretroviral therapy (ART) in low- and middle-income countries (LMIC): a systematic review and meta-analysis. *PloS one*. 2011; **6**(12): e28691.
4. Lawn SD, Badri M, Wood R. Tuberculosis among HIV-infected patients receiving HAART: long term incidence and risk factors in a South African cohort. *AIDS*. 2005; **19**(18): 2109-16.
5. Philip R. An Address on Koch's discovery of the Tubercle Bacillus: Some of its implications and results. *British medical journal*. 1932; **2**(3730): 1-5.
6. Loudon RG, Roberts RM. Droplet expulsion from the respiratory tract. *The American review of respiratory disease*. 1967; **95**(3): 435-42.
7. Loudon RG, Roberts RM. Singing and the dissemination of tuberculosis. *The American review of respiratory disease*. 1968; **98**(2): 297-300.
8. Loudon RG, Spohn SK. Cough frequency and infectivity in patients with pulmonary tuberculosis. *The American review of respiratory disease*. 1969; **99**(1): 109-11.
9. Rouillon A, Perdrizet S, Parrot R. Transmission of tubercle bacilli: The effects of chemotherapy. *Tubercle*. 1976; **57**(4): 275-99.
10. Raviglion M, Sulis G. Tuberculosis 2015: Burden, Challenges and Strategy for Control and Elimination. *Infectious disease reports*. 2016; **8**(2): 6570.
11. Milburn HJ. Primary tuberculosis. *Current opinion in pulmonary medicine*. 2001; **7**(3): 133-41.
12. Dorhoi A, Reece ST, Kaufmann SH. Immunity to intracellular bacteria. Philadelphia: Wolters Kluwer Health, Lippincott Williams & Wilkins; 2012.
13. Dannenberg Jr AM. Macrophage turnover, division and activation within developing, peak and "healed" tuberculous lesions produced in rabbits by BCG. *Tuberculosis*. 2003; **83**(4): 251-60.
14. Cooper AM, Mayer-Barber KD, Sher A. Role of innate cytokines in mycobacterial infection. *Mucosal immunology*. 2011; **4**(3): 252-60.
15. Fletcher HA, Snowden MA, Landry B, Rida W, Satti I, Harris SA, et al. T-cell activation is an immune correlate of risk in BCG vaccinated infants. *Nature communications*. 2016; **7**: 11290.
16. Jambo KC, Banda DH, Afran L, Kankwatira AM, Malamba RD, Allain TJ, et al. Asymptomatic HIV-infected individuals on antiretroviral therapy exhibit impaired lung CD4(+) T-cell responses to mycobacteria. *American journal of respiratory and critical care medicine*. 2014; **190**(8): 938-47.

17. Mariani F, Goletti D, Ciaramella A, Martino A, Colizzi V, Fraziano M. Macrophage response to *Mycobacterium tuberculosis* during HIV infection: relationships between macrophage activation and apoptosis. *Current molecular medicine*. 2001; **1**(2): 209-16.
18. Guirado E, Schlesinger LS. Modeling the *Mycobacterium tuberculosis* Granuloma – the Critical Battlefield in Host Immunity and Disease. *Frontiers in Immunology*. 2013; **4**: 98.
19. Silva Miranda M, Breiman A, Allain S, Deknuydt F, Altare F. The tuberculous granuloma: an unsuccessful host defence mechanism providing a safety shelter for the bacteria? *Clin Dev Immunol*. 2012; **2012**: 139127.
20. Lin PL, Pawar S, Myers A, Pegu A, Fuhrman C, Reinhart TA, et al. Early events in *Mycobacterium tuberculosis* infection in cynomolgus macaques. *Infection and immunity*. 2006; **74**(7): 3790-803.
21. von Reyn CF, Kimambo S, Mtei L, Arbeit RD, Maro I, Bakari M, et al. Disseminated tuberculosis in human immunodeficiency virus infection: ineffective immunity, polyclonal disease and high mortality. *The international journal of tuberculosis and lung disease : the official journal of the International Union against Tuberculosis and Lung Disease*. 2011; **15**(8): 1087-92.
22. Houben RM, Dodd PJ. The Global Burden of Latent Tuberculosis Infection: A Re-estimation Using Mathematical Modelling. *PLoS Medicine*. 2016; **13**(10): e1002152.
23. World Health Organization. Global Tuberculosis Report 2016. Geneva; 2016.
24. GBD 2016 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 328 diseases and injuries for 195 countries, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet*. 2017; **390**(10100): 1211-59.
25. Walter ND, Painter J, Parker M, Lowenthal P, Flood J, Fu Y, et al. Persistent latent tuberculosis reactivation risk in United States immigrants. *American journal of respiratory and critical care medicine*. 2014; **189**(1): 88-95.
26. Lawn SD, Churchyard G. Epidemiology of HIV-associated tuberculosis. *Current opinion in HIV and AIDS*. 2009; **4**(4): 325-33.
27. Jeon CY, Murray MB. Diabetes mellitus increases the risk of active tuberculosis: a systematic review of 13 observational studies. *PLoS Med*. 2008; **5**(7): e152.
28. Dodd PJ, Looker C, Plumb ID, Bond V, Schaap A, Shanaube K, et al. Age- and Sex-Specific Social Contact Patterns and Incidence of *Mycobacterium tuberculosis* Infection. *Am J Epidemiol*. 2016; **183**(2): 156-66.
29. Senkoro M, Kumar AM, Chinnakali P, Mfinanga SG, Egwaga S, Kamara V, et al. Population impact of factors associated with prevalent pulmonary tuberculosis in Tanzania. *The international journal of tuberculosis and lung disease : the official journal of the International Union against Tuberculosis and Lung Disease*. 2016; **20**(10): 1326-33.
30. GBD-2016-Causes-of-Death-Collaborators. Global, regional, and national age-sex specific mortality for 264 causes of death, 1980-2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet*. 2017; **390**(10100): 1151-210.
31. World Health Organisation. Definitions and reporting framework for tuberculosis - 2013 revision (updated December 2014). Geneva: WHO; 2014.

32. World Health Organisation. Antimicrobial resistance: Global report on surveillance. WHO; 2014.
33. Uplekar M, Weil D, Lonnroth K, Jaramillo E, Lienhardt C, Dias HM, et al. WHO's new End TB Strategy. *The Lancet*. 2015; **385**(9979): 1799-801.
34. Bomanji JB, Gupta N, Gulati P, Das CJ. Imaging in tuberculosis. *Cold Spring Harb Perspect Med*. 2015; **5**(6).
35. Shayo GA, Minja LT, Egwaga S, Bakari M, Mugusi FM. Symptom-based screening tool in ruling out active tuberculosis among HIV-infected patients eligible for isoniazid preventive therapy in Tanzania. *Tropical medicine & international health : TM & IH*. 2014; **19**(6): 726-33.
36. National Tuberculosis and Leprosy Programme. Manual for the Management of Tuberculosis and Leprosy. 6th ed. Dar es Salaam: United-Republic-of-Tanzania; 2013.
37. Mtei L, Matee M, Herfort O, Bakari M, Horsburgh CR, Waddell R, et al. High rates of clinical and subclinical tuberculosis among HIV-infected ambulatory subjects in Tanzania. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2005; **40**(10): 1500-7.
38. Lewinsohn DM, Leonard MK, LoBue PA, Cohn DL, Daley CL, Desmond E, et al. Official American Thoracic Society/Infectious Diseases Society of America/Centers for Disease Control and Prevention Clinical Practice Guidelines: Diagnosis of Tuberculosis in Adults and Children. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2017; **64**(2): e1-e33.
39. Global-Laboratory-Initiative. Mycobacteriology laboratory manual. Geneva: WHO Stop TB Partnership. 2014.
40. World Health Organisation. WHO Guidelines Approved by the Guidelines Review Committee. Use of Tuberculosis Interferon-Gamma Release Assays (IGRAs) in Low- and Middle- Income Countries: Policy Statement. Geneva: World Health Organization; 2011.
41. World Health Organisation. WHO Guidelines Approved by the Guidelines Review Committee. Recommendation on 36 Months Isoniazid Preventive Therapy to Adults and Adolescents Living with HIV in Resource-Constrained and High TB- and HIV-Prevalence Settings: 2015 Update. Geneva: WHO; 2015.
42. World Health Organisation. WHO treatment guidelines for drug-resistant tuberculosis 2016 update. Geneva: WHO; 2016.
43. Pathmanathan I, Date A, Coggin WL, Nkengasong J, Piatek AS, Alexander H. Rolling Out Xpert(R) MTB/RIF for TB Detection in HIV-Infected Populations: An Opportunity for Systems Strengthening. *Afr J Lab Med*. 2017; **6**(2).
44. Dheda K, van Zyl Smit R, Badri M, Pai M. T-cell interferon- γ release assays for the rapid immunodiagnosis of tuberculosis: clinical utility in high-burden vs. low-burden settings. *Current opinion in pulmonary medicine*. 2009; **15**(3): 188-200.
45. Banaei N, Gaur RL, Pai M. Interferon Gamma Release Assays for Latent Tuberculosis: What Are the Sources of Variability? *Journal of clinical microbiology*. 2016; **54**(4): 845-50.

46. Clifford V, He Y, Zufferey C, Connell T, Curtis N. Interferon gamma release assays for monitoring the response to treatment for tuberculosis: A systematic review. *Tuberculosis* (Edinburgh, Scotland). 2015; **95**(6): 639-50.
47. Ralph AP, Ardian M, Wiguna A, Maguire GP, Becker NG, Drogumuller G, et al. A simple, valid, numerical score for grading chest x-ray severity in adult smear-positive pulmonary tuberculosis. *Thorax*. 2010; **65**(10): 863-9.
48. Nachiappan AC, Rahbar K, Shi X, Guy ES, Mortani Barbosa EJ, Jr., Shroff GS, et al. Pulmonary Tuberculosis: Role of Radiology in Diagnosis and Management. *Radiographics*. 2017; **37**(1): 52-72.
49. Omri HE, Hascsi Z, Taha R, Szabados L, Sabah HE, Gamiel A, et al. Tubercular Meningitis and Lymphadenitis Mimicking a Relapse of Burkitt's Lymphoma on (18)F-FDG-PET/CT: A Case Report. *Case reports in oncology*. 2015; **8**(2): 226-32.
50. Wang SY, Luo DL, Chen G, Wang SX. 18F-FDG PET/CT Images in a Patient With Primary Chest Wall Tuberculosis Mimicking Malignant Tumor. *Clinical nuclear medicine*. 2016; **41**(4): 323-5.
51. Wang SB, Ji YH, Wu HB, Wang QS, Zhou WL, Lv L, et al. PET/CT for differentiating between tuberculous peritonitis and peritoneal carcinomatosis: The parietal peritoneum. *Medicine*. 2017; **96**(2): e5867.
52. Ankrah AO, van der Werf TS, de Vries EF, Dierckx RA, Sathekge MM, Glaudemans AW. PET/CT imaging of Mycobacterium tuberculosis infection. *Clin Transl Imaging*. 2016; **4**: 131-44.
53. Lefebvre N, Argemi X, Meyer N, Mootien J, Douiri N, Sferrazza-Mandala S, et al. Clinical usefulness of 18F-FDG PET/CT for initial staging and assessment of treatment efficacy in patients with lymph node tuberculosis. *Nucl Med Biol*. 2017; **50**: 17-24.
54. Ye M, Huang J, Wang J, Ren J, Tu J, You W, et al. Multifocal musculoskeletal tuberculosis mimicking multiple bone metastases: a case report. *BMC infectious diseases*. 2016; **16**: 34.
55. Malherbe ST, Shenai S, Ronacher K, Loxton AG, Dolganov G, Kriel M, et al. Persisting positron emission tomography lesion activity and Mycobacterium tuberculosis mRNA after tuberculosis cure. *Nature medicine*. 2016; **22**(10): 1094-100.
56. World Health Organisation. *Treatment of Tuberculosis Guidelines*. 4th ed. Geneva: WHO; 2010.
57. Gualano G, Capone S, Matteelli A, Palmieri F. New Antituberculosis Drugs: From Clinical Trial to Programmatic Use. *Infectious disease reports*. 2016; **8**(2): 6569.
58. Zumla A, Chakaya J, Centis R, D'Ambrosio L, Mwaba P, Bates M, et al. Tuberculosis treatment and management--an update on treatment regimens, trials, new drugs, and adjunct therapies. *The Lancet Respiratory medicine*. 2015; **3**(3): 220-34.
59. Weiss P, Chen W, Cook VJ, Johnston JC. Treatment outcomes from community-based drug resistant tuberculosis treatment programs: a systematic review and meta-analysis. *BMC infectious diseases*. 2014; **14**: 333.
60. Guglielmetti L, Le Du D, Jachym M, Henry B, Martin D, Caumes E, et al. Compassionate use of bedaquiline for the treatment of multidrug-resistant and extensively drug-resistant tuberculosis: interim analysis of a French cohort. *Clinical*

infectious diseases : an official publication of the Infectious Diseases Society of America. 2015; **60**(2): 188-94.

61. Rustomjee R, Zumla A. Delamanid expanded access novel treatment of drug resistant tuberculosis. *Infection and drug resistance*. 2015; **8**: 359-66.
62. Kaufmann SH, Flesch IE. The role of T cell--macrophage interactions in tuberculosis. *Springer Semin Immunopathol*. 1988; **10**(4): 337-58.
63. Kaufmann SH. Protection against tuberculosis: cytokines, T cells, and macrophages. *Ann Rheum Dis*. 2002; **61 Suppl 2**: ii54-8.
64. Gideon HP, Phuah J, Myers AJ, Bryson BD, Rodgers MA, Coleman MT, et al. Variability in tuberculosis granuloma T cell responses exists, but a balance of pro- and anti-inflammatory cytokines is associated with sterilization. *PLoS pathogens*. 2015; **11**(1): e1004603.
65. Dorhoi A, Kaufmann SH. Tumor necrosis factor alpha in mycobacterial infection. *Seminars in immunology*. 2014; **26**(3): 203-9.
66. Khader SA, Partida-Sanchez S, Bell G, Jelley-Gibbs DM, Swain S, Pearl JE, et al. Interleukin 12p40 is required for dendritic cell migration and T cell priming after *Mycobacterium tuberculosis* infection. *The Journal of experimental medicine*. 2006; **203**(7): 1805-15.
67. Drennan MB, Nicolle D, Quesniaux VJ, Jacobs M, Allie N, Mpagi J, et al. Toll-like receptor 2-deficient mice succumb to *Mycobacterium tuberculosis* infection. *The American journal of pathology*. 2004; **164**(1): 49-57.
68. Bafica A, Scanga CA, Feng CG, Leifer C, Cheever A, Sher A. TLR9 regulates Th1 responses and cooperates with TLR2 in mediating optimal resistance to *Mycobacterium tuberculosis*. *The Journal of experimental medicine*. 2005; **202**(12): 1715-24.
69. Banaiee N, Kincaid EZ, Buchwald U, Jacobs WR, Jr., Ernst JD. Potent inhibition of macrophage responses to IFN-gamma by live virulent *Mycobacterium tuberculosis* is independent of mature mycobacterial lipoproteins but dependent on TLR2. *Journal of immunology (Baltimore, Md : 1950)*. 2006; **176**(5): 3019-27.
70. Noss EH, Pai RK, Sellati TJ, Radolf JD, Belisle J, Golenbock DT, et al. Toll-like receptor 2-dependent inhibition of macrophage class II MHC expression and antigen processing by 19-kDa lipoprotein of *Mycobacterium tuberculosis*. *Journal of immunology (Baltimore, Md : 1950)*. 2001; **167**(2): 910-8.
71. Reiley WW, Calayag MD, Wittmer ST, Huntington JL, Pearl JE, Fountain JJ, et al. ESAT-6-specific CD4 T cell responses to aerosol *Mycobacterium tuberculosis* infection are initiated in the mediastinal lymph nodes. *Proceedings of the National Academy of Sciences of the United States of America*. 2008; **105**(31): 10961-6.
72. Mourik BC, Lubberts E, de Steenwinkel JEM, Ottenhoff THM, Leenen PJM. Interactions between Type 1 Interferons and the Th17 Response in Tuberculosis: Lessons Learned from Autoimmune Diseases. *Front Immunol*. 2017; **8**: 294.
73. Su WL, Perng WC, Huang CH, Yang CY, Wu CP, Chen JH. Association of reduced tumor necrosis factor alpha, gamma interferon, and interleukin-1beta (IL-1beta) but increased IL-10 expression with improved chest radiography in patients with pulmonary tuberculosis. *Clinical and vaccine immunology : CVI*. 2010; **17**(2): 223-31.

74. Green AM, Difazio R, Flynn JL. IFN-gamma from CD4 T cells is essential for host survival and enhances CD8 T cell function during Mycobacterium tuberculosis infection. *Journal of immunology (Baltimore, Md : 1950)*. 2013; **190**(1): 270-7.
75. Flynn JL, Chan J, Triebold KJ, Dalton DK, Stewart TA, Bloom BR. An essential role for interferon gamma in resistance to Mycobacterium tuberculosis infection. *The Journal of experimental medicine*. 1993; **178**(6): 2249-54.
76. Mayer-Barber KD, Andrade BB, Barber DL, Hieny S, Feng CG, Caspar P, et al. Innate and adaptive interferons suppress IL-1alpha and IL-1beta production by distinct pulmonary myeloid subsets during Mycobacterium tuberculosis infection. *Immunity*. 2011; **35**(6): 1023-34.
77. Awoniyi DO, Teuchert A, Sutherland JS, Mayanja-Kizza H, Howe R, Mihret A, et al. Evaluation of cytokine responses against novel Mtb antigens as diagnostic markers for TB disease. *The Journal of infection*. 2016; **73**(3): 219-30.
78. Vassalli P. The pathophysiology of tumor necrosis factors. *Annu Rev Immunol*. 1992; **10**: 411-52.
79. O'Garra A, Redford PS, McNab FW, Bloom CI, Wilkinson RJ, Berry MP. The immune response in tuberculosis. *Annu Rev Immunol*. 2013; **31**: 475-527.
80. Chowdhury IH, Ahmed AM, Choudhuri S, Sen A, Hazra A, Pal NK, et al. Alteration of serum inflammatory cytokines in active pulmonary tuberculosis following anti-tuberculosis drug therapy. *Molecular immunology*. 2014; **62**(1): 159-68.
81. Fremont CM, Yermeev V, Nicolle DM, Jacobs M, Quesniaux VF, Ryffel B. Fatal Mycobacterium tuberculosis infection despite adaptive immune response in the absence of MyD88. *The Journal of clinical investigation*. 2004; **114**(12): 1790-9.
82. Wallis RS. Tumour necrosis factor antagonists: structure, function, and tuberculosis risks. *The Lancet Infectious diseases*. 2008; **8**(10): 601-11.
83. Wallis RS, van Vuuren C, Potgieter S. Adalimumab treatment of life-threatening tuberculosis. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2009; **48**(10): 1429-32.
84. Fremont CM, Togbe D, Doz E, Rose S, Vasseur V, Maillet I, et al. IL-1 Receptor-Mediated Signal Is an Essential Component of MyD88-Dependent Innate Response to Mycobacterium tuberculosis Infection. *The Journal of Immunology*. 2007; **179**(2): 1178-89.
85. Juffermans NP, Florquin S, Camoglio L, Verbon A, Kolk AH, Speelman P, et al. Interleukin-1 signaling is essential for host defense during murine pulmonary tuberculosis. *The Journal of infectious diseases*. 2000; **182**(3): 902-8.
86. Mayer-Barber KD, Andrade BB, Oland SD, Amaral EP, Barber DL, Gonzales J, et al. Host-directed therapy of tuberculosis based on interleukin-1 and type I interferon crosstalk. *Nature*. 2014; **511**(7507): 99-103.
87. Khader SA, Bell GK, Pearl JE, Fountain JJ, Rangel-Moreno J, Cilley GE, et al. IL-23 and IL-17 in the establishment of protective pulmonary CD4+ T cell responses after vaccination and during Mycobacterium tuberculosis challenge. *Nature immunology*. 2007; **8**(4): 369-77.

88. Wareham AS, Tree JA, Marsh PD, Butcher PD, Dennis M, Sharpe SA. Evidence for a role for interleukin-17, Th17 cells and iron homeostasis in protective immunity against tuberculosis in cynomolgus macaques. *PloS one*. 2014; **9**(2): e88149.
89. Whittaker E, Nicol M, Zar HJ, Kampmann B. Regulatory T Cells and Pro-inflammatory Responses Predominate in Children with Tuberculosis. *Front Immunol*. 2017; **8**: 448.
90. Redford PS, Murray PJ, O'Garra A. The role of IL-10 in immune regulation during *M. tuberculosis* infection. *Mucosal immunology*. 2011; **4**(3): 261-70.
91. Hussain R, Talat N, Shahid F, Dawood G. Longitudinal tracking of cytokines after acute exposure to tuberculosis: association of distinct cytokine patterns with protection and disease development. *Clinical and vaccine immunology : CVI*. 2007; **14**(12): 1578-86.
92. Tanaka T, Narazaki M, Kishimoto T. IL-6 in inflammation, immunity, and disease. *Cold Spring Harbor perspectives in biology*. 2014; **6**(10): a016295.
93. Kishimoto T. Interleukin-6: discovery of a pleiotropic cytokine. *Arthritis research & therapy*. 2006; **8 Suppl 2**: S2.
94. Nagabhushanam V, Solache A, Ting LM, Escaron CJ, Zhang JY, Ernst JD. Innate inhibition of adaptive immunity: *Mycobacterium tuberculosis*-induced IL-6 inhibits macrophage responses to IFN-gamma. *Journal of immunology (Baltimore, Md : 1950)*. 2003; **171**(9): 4750-7.
95. Zumla A, Rao M, Dodo E, Maeurer M. Potential of immunomodulatory agents as adjunct host-directed therapies for multidrug-resistant tuberculosis. *BMC medicine*. 2016; **14**: 89.
96. Lawn SD, Harries AD, Meintjes G, Getahun H, Havlir DV, Wood R. Reducing deaths from tuberculosis in antiretroviral treatment programmes in sub-Saharan Africa. *Aids*. 2012; **26**(17): 2121-33.
97. Cox JA, Lukande RL, Nelson AM, Mayanja-Kizza H, Colebunders R, Van Marck E, et al. An autopsy study describing causes of death and comparing clinico-pathological findings among hospitalized patients in Kampala, Uganda. *PloS one*. 2012; **7**(3): e33685.
98. Bates M, Mudenda V, Shibemba A, Kaluwaji J, Tembo J, Kabwe M, et al. Burden of tuberculosis at post mortem in inpatients at a tertiary referral centre in sub-Saharan Africa: a prospective descriptive autopsy study. *The Lancet Infectious diseases*. 2015; **15**(5): 544-51.
99. Ansari NA, Kombe AH, Kenyon TA, Hone NM, Tappero JW, Nyirenda ST, et al. Pathology and causes of death in a group of 128 predominantly HIV-positive patients in Botswana, 1997-1998. *The international journal of tuberculosis and lung disease : the official journal of the International Union against Tuberculosis and Lung Disease*. 2002; **6**(1): 55-63.
100. Rana FS, Hawken MP, Mwachari C, Bhatt SM, Abdullah F, Ng'ang'a LW, et al. Autopsy study of HIV-1-positive and HIV-1-negative adult medical patients in Nairobi, Kenya. *Journal of acquired immune deficiency syndromes*. 2000; **24**(1): 23-9.
101. Bryant JM, Harris SR, Parkhill J, Dawson R, Diacon AH, van Helden P, et al. Whole-genome sequencing to establish relapse or re-infection with *Mycobacterium*

- tuberculosis: a retrospective observational study. *The Lancet Respiratory medicine*. 2013; **1**(10): 786-92.
102. Terai C, Kornbluth RS, Pauza CD, Richman DD, Carson DA. Apoptosis as a mechanism of cell death in cultured T lymphoblasts acutely infected with HIV-1. *The Journal of clinical investigation*. 1991; **87**(5): 1710-5.
 103. Chetty S, Govender P, Zupkosky J, Pillay M, Ghebremichael M, Moosa MY, et al. Co-infection with Mycobacterium tuberculosis impairs HIV-Specific CD8+ and CD4+ T cell functionality. *PloS one*. 2015; **10**(3): e0118654.
 104. Sada-Ovalle I, Ocana-Guzman R, Perez-Patrigueon S, Chavez-Galan L, Sierra-Madero J, Torre-Bouscoulet L, et al. Tim-3 blocking rescue macrophage and T cell function against Mycobacterium tuberculosis infection in HIV+ patients. *Journal of the International AIDS Society*. 2015; **18**: 20078.
 105. Espert L, Beaumelle B, Vergne I. Autophagy in Mycobacterium tuberculosis and HIV infections. *Frontiers in cellular and infection microbiology*. 2015; **5**: 49.
 106. Pettit AC, Mendes A, Jenkins C, Napravnik S, Freeman A, Shepherd BE, et al. Timing of Antiretroviral Treatment, Immunovirologic Status, and TB Risk: Implications for Testing and Treatment. *Journal of acquired immune deficiency syndromes*. 2016; **72**(5): 572-8.
 107. Naidoo K, Karim QA, Bhushan A, Yende-Zuma N, McHunu PK, Frohlich J, et al. High rates of tuberculosis in patients accessing HAART in rural South Africa. *Journal of acquired immune deficiency syndromes*. 2014; **65**(4): 438-46.
 108. Bassett IV, Wang B, Chetty S, Giddy J, Losina E, Mazibuko M, et al. Intensive tuberculosis screening for HIV-infected patients starting antiretroviral therapy in Durban, South Africa. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2010; **51**(7): 823-9.
 109. Cox HS, Mbhele S, Mohess N, Whitelaw A, Muller O, Zemanay W, et al. Impact of Xpert MTB/RIF for TB diagnosis in a primary care clinic with high TB and HIV prevalence in South Africa: a pragmatic randomised trial. *PLoS Med*. 2014; **11**(11): e1001760.
 110. Luetkemeyer AF, Firnhaber C, Kendall MA, Wu X, Mazurek GH, Benator DA, et al. Evaluation of Xpert MTB/RIF Versus AFB Smear and Culture to Identify Pulmonary Tuberculosis in Patients With Suspected Tuberculosis From Low and Higher Prevalence Settings. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2016; **62**(9): 1081-8.
 111. Mugusi S, Ngaimisi E, Janabi M, Minzi O, Bakari M, Riedel KD, et al. Liver enzyme abnormalities and associated risk factors in HIV patients on efavirenz-based HAART with or without tuberculosis co-infection in Tanzania. *PloS one*. 2012; **7**(7): e40180.
 112. Manosuthi W, Wiboonchutikul S, Sungkanuparph S. Integrated therapy for HIV and tuberculosis. *AIDS research and therapy*. 2016; **13**: 22.
 113. Belard S, Remppis J, Bootsma S, Janssen S, Kombila DU, Beyeme JO, et al. Tuberculosis Treatment Outcome and Drug Resistance in Lambarene, Gabon: A Prospective Cohort Study. *The American journal of tropical medicine and hygiene*. 2016; **95**(2): 472-80.

114. Bhatt NB, Baudin E, Meggi B, da Silva C, Barrail-Tran A, Furlan V, et al. Nevirapine or efavirenz for tuberculosis and HIV coinfecting patients: exposure and virological failure relationship. *The Journal of antimicrobial chemotherapy*. 2015; **70**(1): 225-32.
115. Kassahun K, McIntosh I, Cui D, Hreniuk D, Merschman S, Lasseter K, et al. Metabolism and disposition in humans of raltegravir (MK-0518), an anti-AIDS drug targeting the human immunodeficiency virus 1 integrase enzyme. *Drug metabolism and disposition: the biological fate of chemicals*. 2007; **35**(9): 1657-63.
116. Sunpath H, Winternheimer P, Cohen S, Tennant I, Chelin N, Gandhi RT, et al. Double-dose lopinavir-ritonavir in combination with rifampicin-based anti-tuberculosis treatment in South Africa. *The international journal of tuberculosis and lung disease : the official journal of the International Union against Tuberculosis and Lung Disease*. 2014; **18**(6): 689-93.
117. Sharma M, Saravolatz LD. Rilpivirine: a new non-nucleoside reverse transcriptase inhibitor. *The Journal of antimicrobial chemotherapy*. 2013; **68**(2): 250-6.
118. Habtewold A, Aklillu E, Makonnen E, Amogne W, Yimer G, Aderaye G, et al. Long-Term Effect of Rifampicin-Based Anti-TB Regimen Coadministration on the Pharmacokinetic Parameters of Efavirenz and 8-Hydroxy-Efavirenz in Ethiopian Patients. *Journal of clinical pharmacology*. 2016.
119. Mariana N, Purwantiastuti, Instiaty, Rusli A. Efavirenz Plasma Concentrations and HIV Viral Load in HIV/AIDS-tuberculosis Infection Patients Treated with Rifampicin. *Acta medica Indonesiana*. 2016; **48**(1): 10-6.
120. Mukonzo JK, Bisaso RK, Ogwal-Okeng J, Gustafsson LL, Owen JS, Aklillu E. CYP2B6 genotype-based efavirenz dose recommendations during rifampicin-based antituberculosis cotreatment for a sub-Saharan Africa population. *Pharmacogenomics*. 2016; **17**(6): 603-13.
121. Naiker S, Connolly C, Wiesner L, Kellerman T, Reddy T, Harries A, et al. Randomized pharmacokinetic evaluation of different rifabutin doses in African HIV- infected tuberculosis patients on lopinavir/ritonavir-based antiretroviral therapy. *BMC pharmacology & toxicology*. 2014; **15**: 61.
122. Mallikaarjun S, Wells C, Petersen C, Paccaly A, Shoaf SE, Patil S, et al. Delamanid Coadministered with Antiretroviral Drugs or Antituberculosis Drugs Shows No Clinically Relevant Drug-Drug Interactions in Healthy Subjects. *Antimicrobial agents and chemotherapy*. 2016; **60**(10): 5976-85.
123. Reniers G, Blom S, Calvert C, Martin-Onraet A, Herbst AJ, Eaton JW, et al. Trends in the burden of HIV mortality after roll-out of antiretroviral therapy in KwaZulu-Natal, South Africa: an observational community cohort study. *Lancet HIV*. 2017; **4**(3): e113-e21.
124. Sirgel FA, Donald PR, Odhiambo J, Githui W, Umapathy KC, Paramasivan CN, et al. A multicentre study of the early bactericidal activity of anti-tuberculosis drugs. *The Journal of antimicrobial chemotherapy*. 2000; **45**(6): 859-70.
125. Jindani A, Aber VR, Edwards EA, Mitchison DA. The early bactericidal activity of drugs in patients with pulmonary tuberculosis. *The American review of respiratory disease*. 1980; **121**(6): 939-49.
126. Mehta JB, Dutt AK, Harvill L, Henry W. Isoniazid preventive therapy for tuberculosis. Are we losing our enthusiasm? *Chest*. 1988; **94**(1): 138-41.

127. Companion Handbook to the WHO Guidelines for the Programmatic Management of Drug-Resistant Tuberculosis. Geneva: World Health Organization 2014.; 2014.
128. Harries AD, Lawn SD, Suthar AB, Granich R. Benefits of combined preventive therapy with co-trimoxazole and isoniazid in adults living with HIV: time to consider a fixed-dose, single tablet coformulation. *The Lancet Infectious diseases*. 2015; **15**(12): 1492-6.
129. Kunkel A, Crawford FW, Shepherd J, Cohen T. Benefits of continuous isoniazid preventive therapy may outweigh resistance risks in a declining TB/HIV co-epidemic. *Aids*. 2016.
130. Danel C, Moh R, Gabillard D, Badje A, Le Carrou J, Ouassa T, et al. A Trial of Early Antiretrovirals and Isoniazid Preventive Therapy in Africa. *The New England journal of medicine*. 2015; **373**(9): 808-22.
131. Samandari T, Agizew TB, Nyirenda S, Tedla Z, Sibanda T, Shang N, et al. 6-month versus 36-month isoniazid preventive treatment for tuberculosis in adults with HIV infection in Botswana: a randomised, double-blind, placebo-controlled trial. *Lancet*. 2011; **377**(9777): 1588-98.
132. Cohen T, Lipsitch M, Walensky RP, Murray M. Beneficial and perverse effects of isoniazid preventive therapy for latent tuberculosis infection in HIV-tuberculosis coinfecting populations. *Proceedings of the National Academy of Sciences of the United States of America*. 2006; **103**(18): 7042-7.
133. Durovni B, Saraceni V, Moulton LH, Pacheco AG, Cavalcante SC, King BS, et al. Effect of improved tuberculosis screening and isoniazid preventive therapy on incidence of tuberculosis and death in patients with HIV in clinics in Rio de Janeiro, Brazil: a stepped wedge, cluster-randomised trial. *The Lancet Infectious diseases*. 2013; **13**(10): 852-8.
134. Ait-Khaled N, Alarcon E, Bissell K, Boillot F, Caminero JA, Chiang CY, et al. Isoniazid preventive therapy for people living with HIV: public health challenges and implementation issues. *The international journal of tuberculosis and lung disease : the official journal of the International Union against Tuberculosis and Lung Disease*. 2009; **13**(8): 927-35.
135. Kabali C, von Reyn CF, Brooks DR, Waddell R, Mtei L, Bakari M, et al. Completion of isoniazid preventive therapy and survival in HIV-infected, TST-positive adults in Tanzania. *The international journal of tuberculosis and lung disease : the official journal of the International Union against Tuberculosis and Lung Disease*. 2011; **15**(11): 1515-21, i.
136. Munseri PJ, Talbot EA, Mtei L, Fordham von Reyn C. Completion of isoniazid preventive therapy among HIV-infected patients in Tanzania. *The international journal of tuberculosis and lung disease : the official journal of the International Union against Tuberculosis and Lung Disease*. 2008; **12**(9): 1037-41.
137. Shayo GA, Moshiri C, Aboud S, Bakari M, Mugusi FM. Acceptability and adherence to Isoniazid preventive therapy in HIV-infected patients clinically screened for latent tuberculosis in Dar es Salaam, Tanzania. *BMC infectious diseases*. 2015; **15**: 368.
138. Takarinda KC, Choto RC, Harries AD, Mutasa-Apollo T, Chakanyuka-Musanhu C. Routine implementation of isoniazid preventive therapy in HIV-infected patients in seven pilot sites in Zimbabwe. *Public Health Action*. 2017; **7**(1): 55-60.

139. Ayele HT, van Mourik MS, Bonten MJ. Predictors of adherence to isoniazid preventive therapy in people living with HIV in Ethiopia. *The international journal of tuberculosis and lung disease : the official journal of the International Union against Tuberculosis and Lung Disease*. 2016; **20**(10): 1342-7.
140. Trinh TT, Han DT, Bloss E, Le TH, Vu TT, Mai AH, et al. Implementation and evaluation of an isoniazid preventive therapy pilot program among HIV-infected patients in Vietnam, 2008-2010. *Transactions of the Royal Society of Tropical Medicine and Hygiene*. 2015; **109**(10): 653-9.
141. LaCourse SM, Deya RW, Graham SM, Masese LN, Jaoko W, Mandaliya KN, et al. Evaluation of the Isoniazid Preventive Therapy Care Cascade Among HIV-Positive Female Sex Workers in Mombasa, Kenya. *Journal of acquired immune deficiency syndromes*. 2017; **76**(1): 74-81.
142. Hoopes AJ, Kammerer JS, Harrington TA, Ijaz K, Armstrong LR. Isoniazid-mono-resistant tuberculosis in the United States, 1993 to 2003. *Archives of internal medicine*. 2008; **168**(18): 1984-92.
143. Wright A, Zignol M, Van Deun A, Falzon D, Gerdes SR, Feldman K, et al. Epidemiology of antituberculosis drug resistance 2002-07: an updated analysis of the Global Project on Anti-Tuberculosis Drug Resistance Surveillance. *Lancet*. 2009; **373**(9678): 1861-73.
144. World Health Organisation. Anti-tuberculosis drug resistance in the world: fourth global report (WHO/HTM/TB/2008.394). Geneva, Switzerland: WHO. 2008.
145. Mills HL, Cohen T, Colijn C. Community-wide isoniazid preventive therapy drives drug-resistant tuberculosis: a model-based analysis. *Science translational medicine*. 2013; **5**(180): 180ra49.
146. Menzies D, Benedetti A, Paydar A, Royce S, Madhukar P, Burman W, et al. Standardized treatment of active tuberculosis in patients with previous treatment and/or with mono-resistance to isoniazid: a systematic review and meta-analysis. *PLoS Med*. 2009; **6**(9): e1000150.
147. Chien JY, Chen YT, Wu SG, Lee JJ, Wang JY, Yu CJ. Treatment outcome of patients with isoniazid mono-resistant tuberculosis. *Clinical microbiology and infection : the official publication of the European Society of Clinical Microbiology and Infectious Diseases*. 2015; **21**(1): 59-68.
148. Nolan C, Goldberg S. Treatment of isoniazid-resistant tuberculosis with isoniazid, rifampin, ethambutol, and pyrazinamide for 6 months. *The International Journal of Tuberculosis and Lung Disease*. 2002; **6**(11): 952-8.
149. Mitchison DA, Nunn AJ. Influence of initial drug resistance on the response to short-course chemotherapy of pulmonary tuberculosis. *The American review of respiratory disease*. 1986; **133**(3): 423-30.
150. Wang TY, Lin SM, Shie SS, Chou PC, Huang CD, Chung FT, et al. Clinical characteristics and treatment outcomes of patients with low- and high-concentration isoniazid-mono-resistant tuberculosis. *PloS one*. 2014; **9**(1): e86316.
151. Villegas L, Otero L, Sterling TR, Huaman MA, Van der Stuyft P, Gotuzzo E, et al. Prevalence, Risk Factors, and Treatment Outcomes of Isoniazid- and Rifampicin-Mono-Resistant Pulmonary Tuberculosis in Lima, Peru. *PloS one*. 2016; **11**(4): e0152933.

152. Yang CS. Advancing host-directed therapy for tuberculosis. *Microb Cell*. 2017; **4**(3): 105-7.
153. Tomioka H, Sano C, Tatano Y. Host-Directed Therapeutics against Mycobacterial Infections. *Curr Pharm Des*. 2017; **23**(18): 2644-56.
154. Skrahin A, Ahmed RK, Ferrara G, Rane L, Poiret T, Isaikina Y, et al. Autologous mesenchymal stromal cell infusion as adjunct treatment in patients with multidrug and extensively drug-resistant tuberculosis: an open-label phase 1 safety trial. *The Lancet Respiratory medicine*. 2014; **2**(2): 108-22.
155. Senkoro M, Mfinanga S, Egwaga S, Mtandu R, Kamara DV, Basra D, et al. Prevalence of pulmonary tuberculosis in adult population of Tanzania: a national survey, 2012. *The international journal of tuberculosis and lung disease : the official journal of the International Union against Tuberculosis and Lung Disease*. 2016; **20**(8): 1014-21.
156. National Tuberculosis and Leprosy Program. The First National Tuberculosis Prevalence Survey in the United Republic of Tanzania Final Report. In: Ministry of Health and Social Welfare, editor. Dar es Salaam: United Republic of Tanzania; 2013.
157. Tanzania Commission for AIDS (TACAIDS), Zanzibar AIDS Commission (ZAC), National Bureau of Statistics (NBS), Office of the Chief Government Statistician (OCGS), and ICF International 2013. Tanzania HIV/AIDS and Malaria Indicator Survey 2011-12. Dar es Salaam, Tanzania: TACAIDS, ZAC, NBS, OCGS, and ICF International.
158. National AIDS Control Program. Costing of the third Health Sector HIV and AIDS Strategic Plan (HSHSP III) 2013-2017. In: Ministry-Of-Health-And-Social-Welfare, editor. Dar es salaam: United-Republic-Of-Tanzania; 2017.
159. Chonde TM, Basra D, Mfinanga SG, Range N, Lwilla F, Shirima RP, et al. National anti-tuberculosis drug resistance study in Tanzania. *The international journal of tuberculosis and lung disease : the official journal of the International Union against Tuberculosis and Lung Disease*. 2010; **14**(8): 967-72.
160. National Tuberculosis and Leprosy Program Xpert® MTB/RIF test Rollout and Implementation Plan. Dar es Salaam: Ministry Of Health And Social Welfare, United Republic of Tanzania.
161. Mkopi A, Range N, Lwilla F, Egwaga S, Schulze A, Geubbels E, et al. Adherence to tuberculosis therapy among patients receiving home-based directly observed treatment: evidence from the United Republic of Tanzania. *PloS one*. 2012; **7**(12): e51828.
162. Egwaga S, Mkopi A, Range N, Haag-Arbenz V, Baraka A, Grewal P, et al. Patient-centred tuberculosis treatment delivery under programmatic conditions in Tanzania: a cohort study. *BMC medicine*. 2009; **7**: 80.
163. Yen YF, Yen MY, Lin YP, Shih HC, Li LH, Chou P, et al. Directly observed therapy reduces tuberculosis-specific mortality: a population-based follow-up study in Taipei, Taiwan. *PloS one*. 2013; **8**(11): e79644.
164. Mugusi SF, Ngaimisi E, Janabi MY, Mugusi FM, Minzi OM, Sasi PG, et al. Risk factors for mortality among HIV-positive patients with and without active tuberculosis in Dar es Salaam, Tanzania. *Antiviral therapy*. 2012; **17**(2): 265-74.

165. Bisson GP, Zetola N, Collman RG. Persistent high mortality in advanced HIV/TB despite appropriate antiretroviral and antitubercular therapy: an emerging challenge. *Curr HIV/AIDS Rep.* 2015; **12**(1): 107-16.
166. Dhasmana DJ, Dheda K, Ravn P, Wilkinson RJ, Meintjes G. Immune reconstitution inflammatory syndrome in HIV-infected patients receiving antiretroviral therapy : pathogenesis, clinical manifestations and management. *Drugs.* 2008; **68**(2): 191-208.
167. Cheng VC, Yam WC, Woo PC, Lau SK, Hung IF, Wong SP, et al. Risk factors for development of paradoxical response during antituberculosis therapy in HIV-negative patients. *European journal of clinical microbiology & infectious diseases : official publication of the European Society of Clinical Microbiology.* 2003; **22**(10): 597-602.
168. Meintjes G, Rabie H, Wilkinson RJ, Cotton MF. Tuberculosis-associated immune reconstitution inflammatory syndrome and unmasking of tuberculosis by antiretroviral therapy. *Clin Chest Med.* 2009; **30**(4): 797-810.
169. Dannenberg AM, Dey B. Perspectives for Developing New Tuberculosis Vaccines Derived from the Pathogenesis of Tuberculosis: I. Basic Principles, II. Preclinical Testing, and III. Clinical Testing. *Vaccines.* 2013; **1**(1): 58-76.
170. Fletcher JM, Vukmanovic-Stejić M, Dunne PJ, Birch KE, Cook JE, Jackson SE, et al. Cytomegalovirus-specific CD4⁺ T cells in healthy carriers are continuously driven to replicative exhaustion. *Journal of immunology (Baltimore, Md : 1950).* 2005; **175**(12): 8218-25.
171. Ellertsen LK, Storla DG, Diep LM, Brokstad KA, Wiker HG, Hetland G. Allergic sensitisation in tuberculosis patients at the time of diagnosis and following chemotherapy. *BMC infectious diseases.* 2009; **9**: 100.
172. Djoba Siawaya JF, Beyers N, van Helden P, Walzl G. Differential cytokine secretion and early treatment response in patients with pulmonary tuberculosis. *Clinical and experimental immunology.* 2009; **156**(1): 69-77.
173. Ladel CH, Blum C, Dreher A, Reifenberg K, Kopf M, Kaufmann SH. Lethal tuberculosis in interleukin-6-deficient mutant mice. *Infection and immunity.* 1997; **65**(11): 4843-9.
174. Tang S, Cui H, Yao L, Hao X, Shen Y, Fan L, et al. Increased cytokines response in patients with tuberculosis complicated with chronic obstructive pulmonary disease. *PloS one.* 2013; **8**(4): e62385.
175. Casarini M, Ameglio F, Alemanno L, Zangrilli P, Mattia P, Paone G, et al. Cytokine levels correlate with a radiologic score in active pulmonary tuberculosis. *American journal of respiratory and critical care medicine.* 1999; **159**(1): 143-8.
176. Ravimohan S, Tamuhla N, Steenhoff AP, Letlhogile R, Nfanyana K, Bellamy SL, et al. Immunological profiling of tuberculosis-associated immune reconstitution inflammatory syndrome and non-immune reconstitution inflammatory syndrome death in HIV-infected adults with pulmonary tuberculosis starting antiretroviral therapy: a prospective observational cohort study. *The Lancet Infectious diseases.* 2015.
177. Ravimohan S, Tamuhla N, Nfanyana K, Steenhoff AP, Letlhogile R, Frank I, et al. Robust reconstitution of TB-specific polyfunctional CD4⁺ T-cell responses and rising systemic IL-6 in paradoxical TB-IRIS. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America.* 2015.

178. Verbon A, Juffermans N, Van Deventer SJ, Speelman P, Van Deutekom H, Van Der Poll T. Serum concentrations of cytokines in patients with active tuberculosis (TB) and after treatment. *Clinical and experimental immunology*. 1999; **115**(1): 110-3.
179. Zumla A, Rao M, Parida SK, Keshavjee S, Cassell G, Wallis R, et al. Inflammation and tuberculosis: host-directed therapies. *J Intern Med*. 2015; **277**(4): 373-87.
180. Hnizdo E, Singh T, Churchyard G. Chronic pulmonary function impairment caused by initial and recurrent pulmonary tuberculosis following treatment. *Thorax*. 2000; **55**(1): 32-8.
181. de Valliere S, Barker RD. Residual lung damage after completion of treatment for multidrug-resistant tuberculosis. *The international journal of tuberculosis and lung disease : the official journal of the International Union against Tuberculosis and Lung Disease*. 2004; **8**(6): 767-71.
182. Rossi JF, Lu ZY, Jourdan M, Klein B. Interleukin-6 as a therapeutic target. *Clinical cancer research : an official journal of the American Association for Cancer Research*. 2015; **21**(6): 1248-57.
183. ClinicalTrials.gov. Siltuximab therapy in clinical trials. [Electronic] 2016 [cited 2016 11.10.2016]; Available from: <https://clinicaltrials.gov/ct2/results?term=siltuximab&Search=Search>
184. ClinicalTrials.gov. Tocilizumab therapy in clinical trials. [Electronic] 2016 [cited 2016 11.10.2016]; Available from: <https://clinicaltrials.gov/ct2/results?term=tocilizumab&Search=Search>
185. Okada M, Kita Y, Kanamaru N, Hashimoto S, Uchiyama Y, Mihara M, et al. Anti-IL-6 receptor antibody causes less promotion of tuberculosis infection than anti-TNF-alpha antibody in mice. *Clinical & developmental immunology*. 2011; **2011**: 404929.
186. Singh PP, Goyal A. Interleukin-6: a potent biomarker of mycobacterial infection. *SpringerPlus*. 2013; **2**: 686.
187. Leal IS, Smedegard B, Andersen P, Appelberg R. Interleukin-6 and interleukin-12 participate in induction of a type 1 protective T-cell response during vaccination with a tuberculosis subunit vaccine. *Infection and immunity*. 1999; **67**(11): 5747-54.